PCT

cited in the European Search Report of EP0092 35565 Your Ref.: PF 354PCT3EP

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

C12N 15/12, 15/85, 5/10, C07K 14/705, 16/28, C12Q 1/68, A61K 38/17, 48/00

(11) International Publication Number:

WO 98/31799

(43) International Publication Date:

23 July 1998 (23.07.98)

(21) International Application Number:

PCT/US98/00959

A3

(22) International Filing Date:

21 January 1998 (21.01.98)

(30) Priority Data:

60/034,204 60/034,205 21 January 1997 (21.01.97) 21 January 1997 (21.01.97)

US US

(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US). GENTZ, Reiner, L. [DE/US]; 13404 Fairland Park Drive, Silver Spring, MD 20904 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US).
- (74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(88) Date of publication of the international search report:
5 November 1998 (05.11.98)

(54) Title: POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

(57) Abstract

Receptor polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing receptor polypeptides and polynucleotides in the design of protocols for the treatment of diseases and diagnostic assays for such conditions.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	1E	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	15	Iceland	MW	Malawi	US	United States of America
CA	Canada	Γſ	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan .	NE	Niger	VN	Vict Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal	٠.	
CU	Cuba	KZ	Kazakstan	RO	Romania		
cz	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	и	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

WO 98/31799 PCT/US98/00959

POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

FIELD OF INVENTION

5

10

15

20

25

30

35

This invention relates to newly identified polynucleotides and the polypeptides encoded by them, the use of such polynucleotides and polypeptides, and their production. More particularly, the polynucleotides and polypeptides of the present invention relate to specific receptor families described in the specification and known in the art. The invention also relates to inhibiting or activating the action of such polynucleotides and polypeptides.

BACKGROUND OF THE INVENTION

Receptor proteins are found on the membrane of the cells and are generally involved in signal transduction. There are many types of receptor proteins, and for convenience, these proteins are grouped in families based on similarity in structure and function.

For example, the TM4SF superfamily of cell surface proteins, also known as the tetraspan receptor superfamily, is comprised of at least seventeen individual gene products (these include CD9, CD20, CD37, CD53, CD63, CD81, CD82, A15, CO-029, Sm23, RDS, Uro B, Uro A, SAS, Rom-1, PETA3, and YKK8). The TM4SF superfamily is the second largest group in the CD antigen superfamily. Each member of the TM4SF superfamily can be characterized by several putative physical features including four highly conserved transmembrane domains, two divergent extracellular loops, and two short and highly divergent cytoplasmic tails. Expression patterns for members of the TM4SF superfamily tend to be rather broad and can vary widely between members. The functional roles of TM4SF superfamily members are primarily associated with signal transduction events and pathways, but also include cell adhesion in platelets and other lymphocytic and non-lymphocytic cell lines, as well as cell motility, proliferation, and metastasis. In addition, recent evidence suggests that a subset of the members of the TM4SF superfamily may function as potassium channel molecules.

One member of the TM4SF family, CD20, is a four membrane spanning domain cell surface phosphoprotein expressed exclusively on B lymphocytes. Although the precise functional role of CD20 has yet to be determined, it is thought to function primarily as a receptor during B-cell activation. Furthermore, a large number of experimental observations suggest several additional speculative roles for the CD20 molecule. For example, CD20-specific immunoprecipitation of biochemically cross-linked plasma membrane proteins suggests that CD20 assumes a multimeric structural

conformation characteristic of other previously described membrane channel proteins. Further experimentation has revealed that expression of exogenous CD20 on the cell surface specifically increases Ca²⁺ conductance across the plasma membrane. Together, these results suggest that CD20 complexes may function as B-cell specific Ca²⁺ ion channels. In addition, monoclonal antibodies raised against CD20 have been used to stimulate resting B-cells to transition out of the G0/G1 segment of the cell cycle. It has also been demonstrated that CD20 is associated with both serine and tyrosine kinases and, more specifically, that CD20 is associated, although not directly, with the Src family of tyrosine kinases including p56/53lyn, p56lck, and p59fyn.

PCT/US98/00959

不会一切以外的人 医精神的 医阿特克斯氏氏病病 化化丁基乙基甲基甲二基甲基乙基

A second example of a receptor subfamily, called sialoadhesin molecules, belongs to the Ig superfamily of receptor-like molecules. The more than 100 members of the Ig superfamily are generally considered to engage in specific cell-cell interactions through which intercellular communication may occur. In addition to classical protein-protein interactions, intercellular communication may also be mediated through protein-carbohydrate interactions. In fact, all members of the sialoadhesin family of the Ig superfamily are capable of mediating protein-sialic acid binding interactions. To date, only a small number of proteins have been assigned to the sialoadhesin family including sialoadhesin, CD33, CD22, the myelin-associated glycoprotein (MAG), and the Schwann cell myelin protein (SMP). Each of these proteins is expressed in a restricted subset of cell types. For example, CD22 and CD33 are expressed exclusively by B-lymphocytes and cells of the myelomonocytic lineage, respectively.

Similarly, galectins are a family of the lectin superfamily of carbohydrate-binding proteins which have a high affinity for b-galactoside sugars. Although a large number of glycoproteins containing b-galactoside sugars are produced by the cell, only a few will bind to known galectins *in vitro*. Such apparent binding specificity suggests a highly specific functional role for the galectins. Galectin 1 (conventionally termed *LGALS1* for lectin, galactoside-binding, soluble -1) is thought to specifically bind laminin, a highly polylactosaminated cellular glycoprotein, as well as the highly polylactosaminated lysosome-associated membrane proteins (LAMPs). Galectin 1 has also been shown to bind specifically to a lactosamine-containing glycolipid found on olfactory neurons and to integrin a₇b₁ on skeletal muscle cells. Galectin 3 has also been observed to bind specifically to laminin, immunoglobulin E and its receptor, and bacterial lipopolysaccharides.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin 1 appears to inhibit cell adhesion of skeletal muscle presumably by galectin 1-mediated disruption of laminin-integrin a_7b_1 interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined

5

10

15

20

25 -

30

presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bc12, respectively.

A relatively new family of cell-surface proteins has been identified and termed the Ly6 superfamily. The members of this family include murine and human SCA-2, rat Ly-6 (also termed ThB), human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)], and E48 antigen. The determination of an initial functional role for SCA-2 may lie in an analysis of its expression profile with regard to the complex process of hematopoiesis. SCA-2 is highly expressed in early thymic precusor cells. In turn, progeny of the intrathymic precusor population continue to express SCA-2, but only until the point of transition occurs from blast cell to small cell. Further experimental evidence demonstrates that mature thymocytes and peripheral T-cells do not express detectable levels of SCA-2, whereas mature, peripheral B-cells do continue to express SCA-2. As a result, it seems very likely that SCA-2 plays an important role in thymocyte maturation and differentiation. A plausible explanation for this functional hypothesis is that SCA-2 may act as a receptor for a unknown cytokine which regulates thymocyte maturation and differentiation.

In addition, CD59 is a recently identified integral membrane protein which appears to be involved in the regulation of complement. Recent studies show that the CD59 antigen may prevent damage from complement C5b-9 and protect astrocytes during inflammatory and infectious disorders of the nervous system. Expression of recombinant human CD59 on porcine donor organs have been shown to prevent complement-mediated lysis and activation of endothelial cells that leads to hyperacute rejection. Recently, researchers at Alexion Pharmaceuticals (New Haven, CT) reported on the production of transgenic pigs which expressed human CD59. In these animals, xenogeneic organs were resistant to hyperacute rejection. (Fodor, et al., "Expression of a functional human complement inhibitor in a transgenic pig as a model for the prevention of xenogeneic hyperacute organ rejection," Proc. Natl. Acad. Sci., 91:1153-11157 (1994).) The same company also reported that expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria (PNH) B-

5

10

15

20

25

30

WO 98/31799 4 PCT/US98/00959

cells confers resistance to human complement. (Rother et al., "Expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria B-cells confers resistance to human complement," Blood, 84:2604-2611 (1994).) PNH is an acquired hematopoietic disorder characterized by complement-mediated hemolytic anemia, pancytopenia, and venous thrombosis. It is thought that retroviral gene therapy with this molecule could provide a treatment for PNH patients.

A final Ly6 superfamily member, the E48 antigen, is involved in intercellular adhesion between keratinocyte cells of the squamous epithelium. Such keratinocytes are attached to adjoining cells by large numbers of desmosomes, which are thought to play a role in the transition of transformed keratinocytes to metastatic tumor cells. Treatment with a monoclonal antibody raised against the E48 antigen has been successful in the eradication of residual, postoperative squamous cell carcinoma cells of the upper aerodigestive tract in several *in vivo* models and, to some degree, in humans. (van Dongen, et al., "Progress in radioimmunotherapy of head and neck cancer," Oncol. Rep. 1:259-264 (1994).) The gene encoding the E48 antigen has been mapped to the q24-qter region of human chromosome 8. Interestingly, a number of human diseases have been mapped to this region of chromosome 8 including Langer-Giedion syndrome, brachio-otorhinolaryngeal syndrome, trichorhinolaryngeal syndrome, and epidermolysis bullosa simplex.

and the state of the state of the state of

A further example of a receptor family includes the prohibitin receptors. The prohibitin gene product is expressed in a wide variety of tissues and has been implicated as a component of a number of anti-proliferative mechanisms. The prohibitin gene encodes a 30 kD postsynthetically modified polypeptide located primarily in the mitochondria, but also may be associated with the IgM receptor on the B-cell plasma membrane. The protein functionally inhibits DNA synthesis and entry into S phase of the cell cycle by an unknown mechanism. Interestingly, although the prohibitin gene product is hypothesized to be involved in the maintenance of senescence and the prevention of cancer, one study found that, although somatic mutations in the prohibitin gene were present in a small number of breast cancers, no mutations were identified in any other breast, ovary, liver, and lung cancers examined. (Sato et al., Genomics 17:762-764 (1993).) However, the prohibitin gene has been mapped to human chromosome 17q12-21, the same region thought to contain the gene involved in sporadic breast cancer. Furthermore, DNA sequence analysis of the prohibitin gene identified somatic mutation in 4 of 23 cases of sporadic breast cancer examined. Thus, prohibitin family members may be involved in the development of cancer.

Moreover, the EGFR family of plasma membrane proteins are an integral component of normal cellular proliferation and in the pathogenesis of the cancerous state. The family is relatively small and includes the EGFR, c-erbB-2, c-erbB-3, and others. Various cancers are correlated with aberrant expression of one or more of these

5

10

15

20

25

30

10

15

20

25

30

35

genes. A number of ligands have been identified which bind to the EGFR-like receptors listed above including TGF-a, heparin-binding EGF, amphiregulin, criptoregulin, heregulin, and others. A large fraction of adenocarcinomas examined to date, especially those of the breast, colon, and pancreas, are typified by the amplification or overexpression of the c-erbB-2 gene. EGF, or an analogous ligand, initiates the cellular growth factor response by binding to the EGFR, or EGFR-related, receptor. Following the binding event, the receptor molecule dimerizes activating its intracellular tyrosine kinase domain. This event results in the phosphorylation of specific tyrosine residues near the carboxy terminus of the receptor. The diversity of signals able to be transduced through the relatively small number of EGFR-related receptor molecules is amplified considerably by the recent finding that EGFR-like receptor molecules can function when dimerized with other EGFR family members forming heterodimers.

Members of the EGFR-related family of integral membrane proteins have been implicated in the pathogenesis of a number of human disease-states. For example, a mutation in the EGFR itself appears to play an important role in the development of glioblastomas. (Sang et al., J. Neurosurg 82:841-846 (1995).) The EGFR gene is amplified or overexpressed in the majority of primary human glioblastomas. Although not conferring a distinct advantage on cell growth, an increase in EGFR expression was found to confer an increase in the ability of glioma cells to maintain anchorage-independent growth in soft agar especially in response to EGF and retinoic acid. Anchorage-independent growth in vitro correlates highly with tumorigenicity in vivo, therefore, it is likely that cells which express abnormally high levels of EGFR in human glioblastoma cells may be involved in the high potential for these cells to cause tumors in vivo.

Moreover, overexpression or amplification of c-erbB-2 has been reported to be involved in a high number adenocarcinomas, particularly of the breast, colon, and pancreas, and in a small proportion of ovarian carcinomas.

Thus, there is a clear need for identifying and exploiting novel members of the receptor families, such as those described above. Although structurally related, these receptors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such receptors may prove useful as therapeutics in an antitumor, diagnostic, or other capacity. Furthermore, receptors described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

SUMMARY OF THE INVENTION

In one aspect, the invention relates to receptor polypeptides and polynucleotides, as well as the methods for their production. Another aspect of the invention relates to methods for using such receptor polypeptides and polynucleotides.

5 Such uses include the treatment of the specified diseases, among others. In still another aspect, the invention relates to methods to identify agonists and antagonists using the materials provided by the invention, and treating conditions associated with receptor imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate receptor activity or levels.

ъ

DESCRIPTION OF THE INVENTION Definitions

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Receptor" refers, among others, to a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:Y, or an allelic variant thereof.

"Receptor Activity" or "Biological Activity of the Receptor" refers to the metabolic or physiologic function of said receptor including similar activities or improved activities or these activities with decreased undesirable side-effects. Also included are antigenic and immunogenic activities of said receptor.

"Receptor gene" refers to a polynucleotide comprising the nucleotide sequence set forth in SEQ ID NO:X or allelic variants thereof and/or their complements.

"SEQ ID NO:X" comprises all or a substantial portion of the polynucleotide encoding each receptor of the invention. The value X for the nucleotide sequence is an integer specified in Table 1. This nucleotide sequence was translated into the receptor polypeptide identified in Table 1 as "SEQ ID NO:Y," where the value of Y for each receptor polypeptide is an integer defined in Table 1.

The invention further provides a composition of matter comprising a nucleic acid molecule which comprises a human cDNA clone identified by a cDNA Clone ID (Identifier) in Table 1, which DNA molecule is contained in the material deposited with the American Type Culture Collection ("ATCC") and given the ATCC Deposit Number shown in Table 1 for that cDNA clone. The ATCC is located at American Type Culture Collection (ATCC), 12301 Park Lawn Drive, Rockville, Maryland 20852, USA. The deposit has been made under the terms of the Budapest Treaty on the international recognition of the deposit of micro-organisms for purposes of patent procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposit is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that

15

20

25

30

required under 35 U.S.C. §112. The nucleotide sequence of the polynucleotides contained in the deposited material, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can

5

10

15

20

25

30

WO 98/31799 ⁸ PCT/US98/00959

occur anywhere in a polypeptide, including the peptide backbone, the amino acid sidechains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphotidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter et al., "Analysis for protein modifications and nonprotein cofactors", Meth Enzymol (1990) 182:626-646 and Rattan et al., "Protein Synthesis: Posttranslational Modifications and Aging", Ann NY Acad Sci (1992) 663:48-62.)

"Variant" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to

5

10

15

20

25

30

10

15

20

occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" per se has an art-recognized meaning and can be calculated using published techniques. (See, e.g.: COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; SEOUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991.) While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans. (Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073.) Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCS program package (Devereux, J., et al., Nucleic Acids Research (1984) 12(1):387), BLASTP, BLASTN, FASTA (Atschul, S.F. et al., J Molec Biol (1990) 215:403.)

25

30

35

As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO:X is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: X. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5 ' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

Similarly, by a polypeptide having an amino acid sequence having at least, for example, 95% "identity" to a reference amino acid sequence of SEQ ID NO:Y is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO:Y. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

15

20

10

5

Polypeptides of the Invention

In one aspect, the present invention relates to receptor polypeptides (or receptor proteins). The receptor polypeptides include the polypeptide of SEQ ID NO:Y; as well as polypeptides comprising the amino acid sequence of SEQ ID NO:Y; and polypeptides comprising the amino acid sequence which have at least 80% identity to that of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y. Furthermore, those with at least 97-99% identity to SEQ ID NO:Y are highly preferred. Also included within receptor polypeptides are polypeptides having the amino acid sequence which have at least 80% identity to the polypeptide having the amino acid sequence of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y. Furthermore, those with at least 97-99% are highly preferred. Preferably receptor polypeptides exhibit at least one biological activity of the receptor.

30

The receptor polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, prosequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

35

Fragments of the receptor polypeptides are also included in the invention. A "fragment" is a polypeptide having an amino acid sequence that entirely is the same as part, but not all, of the amino acid sequence of the aforementioned receptor polypeptides. As with receptor polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most

10

15

20

25

30

35

preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101 to the end of receptor polypeptide. In this context "about" includes the particularly recited ranges larger or smaller by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of receptor polypeptides, except for deletion of a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus.

Also preferred are fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. The "domains" of each receptor polypeptide are illustrated in the Figures. The Figures compare SEQ ID NO:Y to the closest know homologue. Identical amino acids shared between the two polypeptides are shaded, while conservative amino acid changes are boxed. By examining the regions or amino acids shaded and/or boxed, the skilled artisan can readily identify conserved domains between the two polypeptides. The amino acids sequences of SEQ ID NO:Y falling within these conserved domains are "fragments" and are specifically contemplated by the present invention. Especially preferred is the extracellular domains of a receptor of the invention. Soluble extracellular domains have antagonist activity mediated by competition with a receptor ligand.

Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate receptor activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that are antigenic or immunogenic in an animal, especially in a human.

Preferably, all of these polypeptide fragments retain a biological activity of the receptor, including antigenic activity. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents by conservative amino acid substitutions -- i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination.

The receptor polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

12

Polynucleotides of the Invention

5

10

15

20

30

35

Another aspect of the invention relates to receptor polynucleotides. Receptor polynucleotides include isolated polynucleotides which encode the receptor polypeptides and fragments, and polynucleotides closely related thereto. More specifically, a receptor polynucleotide of the invention includes a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:X encoding a receptor polypeptide of SEQ ID NO:Y, and polynucleotide having the particular sequence of SEQ ID NO:X.

Receptor polynucleotides further include a polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y, and a polynucleotide comprising a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. Also included under receptor polynucleotides are a nucleotide sequence which has sufficient identity to a nucleotide sequence contained in SEQ ID NO:X, or contained in the cDNA insert in the plasmid deposited with ATCC, to hybridize under conditions useable for amplification or for use as a probe or marker. Moreover, the receptor polynucleotide includes a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC, and a nucleotide sequence comprising at least 15 contiguous nucleotides of such cDNA insert. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. The invention also provides polynucleotides which are complementary to all the above receptor polynucleotides.

The receptors of the invention are structurally related to other proteins of specified receptor families, as shown by the results in the Figures. The cDNA sequence of SEQ ID NO:X encodes a polypeptide as described in Table 1 as SEQ ID NO:Y. Because the receptor polypeptides contain domains similar in structure to other receptor family members, the receptors of the present invention are expected to have,

inter alia, similar biological functions/properties to their homologous polypeptides and polynucleotides, and their utility is obvious to anyone skilled in the art.

Table 1

5

Clone ID	SEQ	SEQ	ATCC	ATCC	Receptor	Homology
Name	ID NO:X	ID NO:Y	Deposit No.	Deposit Date	Family	,
HMACR70			209054	05/16/97	Ig	Sialoadhesin
	1	18	#####	01/21/98		OB-1
HTEDK48			209054	05/16/97	TM4SF	MRC-OX44
				<u> </u>		PETA-3
1-1849 bp	2					
160-900 bp	3	19				
HTPED39			209054	05/16/97	TM4SF	NAG-2
HPWAE25	4	20	#####	1/21/98		TALLA-1
HTPEF86	5	21	209053	05/16/97	TM4SF	CD20
						B1 Antigen
HSBBF02	6	22	209054	05/16/97	TM4SF	TALLA-1
HLTAH80	7	23	97242	08/02/95	TM4SF	TALLA-1
	L		209054	05/16/97		
HTPBA27	8	24	97242	08/02/95	TM4SF	NAG-2
			209054	05/16/97		675
HAIDQ59			209054	05/16/97	TM4SF	CD9
						Antigen
5' Sequence	9	25				
3' Sequence	10				50.7.405	705754.0
HHFEK40	11	26	209054	05/16/97	TM4SF	PETA-3
HGBGV89	12	27	209125	06/09/97	TM4SF	L6H
			209054	05/16/97		
HUVBB80	13	28	209054	05/16/97	TM4SF	L6
HJACE54	14	29	209053	05/16/97	Lectin	Galectin-3
				ł		Galectin-5
				254165	<u> </u>	Galectin-8
HROAD63	15	30	209053	05/16/97	Ly6	E48 splice
					<u> </u>	variant
HMWGS46	16	31	209053	05/16/97	Prohibitin	BAP-37
HNFGW06	17	32	209053	05/16/97	EGFR	EGFR

The novel full-length cDNA clone designated HMACR70 may be a member of the sialoadhesin family of the Ig superfamily of receptor-like molecules and a CD33 homologue. HMACR70 contains a 1497 nucleotide cDNA insert encoding a 315 amino acid ORF and was cloned from a GM-CSF-treated human macrophage cDNA library. The only additional cDNA libraries in the HGS database which include this clone are human eosinophils and possibly human gall bladder. A BLAST analysis of the amino acid sequence of HMACR70 demonstrates that this clone exhibits approximately 50% identity and 69% similarity over a 300 amino acids stretch of a gene termed human

10

differentiation antigen, and 38% identity and 62% similarity of the human myelin-associated glycoprotein precursor CD33 gene.

A more recent BLAST analysis confirms HMACR70's designation as a sialoadhesin family member. HMACR70 is homologous to two recently identified sialoadhesin family members, human OB binding protein (OB) 1 and 2. (See, Genbank Accession No. U71382; see Figure 1.) It is thought that OB-1 and OB-2 may bind leptin. Thus, HMACR70, as a sialoadhesin family member, may act to attenuate or even amplify intercellular routes of communication, including binding to leptin or modulating the activity of immune cells, such as macrophages. Clearly, any diseases affected by these processes could be treated by the polypeptide or fragment of HMACR70.

The full-length nucleotide sequences of ten novel human cDNA clones which potentially belong to the TM4SF superfamily are disclosed in the table above and will be addressed sequentially.

The cDNA clone HTEDK48 contains a 1849 nucleotide cDNA insert encoding a 245 amino acid ORF that was cloned from a human testes cDNA library. The coding sequence of HTEDK48 (SEQ ID NO: 3) may be fused to other human proteins, such as 3-hydroxyacyl-CoA dehydrogenase. BLAST analysis of the amino acid sequence of HTEDK48 demonstrates that this clone exhibits approximately 30% identity and 51% similarity over a 245 amino acid stretch of the CD82 molecule. Recent studies have shown that CD82 can associate with CD4 or CD8 and deliver costimulatory signals for the TCR/CD3 pathway. CD82 has also been found to be involved in syncytium formation in HTLV-I-infected T-cells. And finally, in a recently published study in which the expression of the CD82 gene by tumors of the lung was examined retrospectively, it was reported that CD82 may be linked to the suppression of tumor metastasis of prostate cancer. The study also reported that decreased CD82 expression may be involved in malignant progression of such cancers. Thus, HTEDK48 may also be involved in the development of cancer.

The state of the s

A more recent BLAST analysis shows that HTEDK48 is homologous the rat leukocyte antigen, MRC OX-44, and the platelet endothelial tetraspan antigen -3 (PETA-3). (See Figure 2X.) MRC OX-44, a member of a new family of cell surface proteins, appears to be involved in growth regulation. (See, Bellacosa, A., et al., "The Rat Leukocyte antigen MRC OX-44 is a Member of a New Family of Cell Surface Proteins which Appear to be Involved in Growth Regulation," Mol. Cell. Bio. 11: 2864-2872 (1991).) Similarly, PETA-3 has been located to platelet endothelial cells, and an anti-PETA-3 antigen monoclonal antibody can stimulate platelet aggregation and mediator release. (See, Fitter, S., "Molecular Cloning of cDNA Encoding a Novel Platelet-Endothelial Cell Tetra-Span Antigen, PETA-3," Blood, 86(4):1348-1355 (1995).) Thus, HTEDK48 may function similar to MRC OX-44 or PETA-3 to affect

5

10

15

20

25

30

WO 98/31799 15 PCT/US98/00959

growth of blood cells. Administering polypeptides or fragments of HTEDK48 may be an effective treatment of blood disorders.

The cDNA clone **HPWAE25** contains a 1288 nucleotide cDNA insert encoding a 273 amino acid ORF that was cloned from a human pancreas tumor cDNA library, while clone **HTPED39** represents a truncated cDNA sequence. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including keratinocytes, ulcerative colitis, striatum depression, lymph node breast cancer, ovarian cancer, stage B2 prostate cancer, kidney medulla, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human cell lines including U937, MM96, WM115, and MDAMB231. A BLAST analysis of the amino acid sequence of HTPED39 demonstrates that this clone exhibits approximately 35% identity and 50% similarity over the entire length of the CD37 molecule. The CD37 antigen is expressed on B cells and on a subpopulation of T cells, but not on pre-B or plasma cells. It has been reported that CD37 expression is downregulated in conjunction with B-cell activation, suggesting that CD37 may be involved in the processes which dictate the activation state of the B-cell.

Moreover, HPWAE25 is also homologous to recently identified TM4SF members, NAG-2 and TALLA-1. (See Figure 3.) NAG-2 is thought to complex with integrins and other TM4SF proteins, while TALLA-1 is a highly specific marker of T-cell acute lymphoblastic leukemia and neuroblastoma. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that Complexs with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997); Takagi, S., "Identification of a Higly Specific Surface Marker of T-cell Acute Lymphoblastic Leukemia and Neuroblastoma as a New Member of the Transmembrane 4 Superfamily," Int. J. Cancer 61(5):706-715 (1995).) Thus, HPWAE25 may be involved the development of cancer, particularly leukemia, lymphoma, and neuroblastoma. HPWAE25 may be used as an effective treatment of these cancers, as well as a diagnostic marker.

A subfamily of TM4SF receptors include CD20 proteins. A CD20-like cDNA clone was obtained from a human pancreas tumor cDNA library and contains a 1236 nucleotide insert which encodes a 250 amino acid ORF. A BLAST analysis of the deduced amino acid sequence of HTPEF86 exhibits approximately 41% identity and 61% similarity to the CD20 gene, also known as B1 antigen. (See Figure 4.) Expression of this gene is detected in only two additional HGS human cDNA libraries; amygdala depression and 9 week early stage human. Although the precise functional role of CD20 has yet to be determined, it is clear that CD20 plays a key role in the regulation of B-cell activation. Based primarily on sequence identity, the novel CD20-like molecule presented herein may also be involved in cell cycle activation. Potential therapeutic and/or diagnostic applications for HTPEF86 may include such clinical

5

10

15

20

25

30

WO 98/31799 16 PCT/US98/00959

presentations as juvenile rheumatoid arthritis, Graves' Disease, and a number of B-cell lymphomas or other lymphoid tumors.

The clone **HSBBF02** contains a 1115 nucleotide cDNA insert encoding a 245 amino acid ORF and was cloned from an HSC 172 cell line cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including brain amygdala depression, endothelial cells, fetal liver and heart, osteoblasts, testes, and others. A BLAST analysis of the amino acid sequence of HSBBF02 demonstrates that this clone exhibits approximately 64% identity and 80% similarity with the A15 molecule over a 131 amino acid stretch (A15 is composed of 244 amino acids). A more recent BLAST search shows that HSBBF02 is similar to the TALLA-1 protein and may in fact be a closely related family member. (See Figure 5.)

In addition, a second cDNA clone, designated HLTAH80, exhibits sequence similarity to the A15 molecule and TALLA-1. (See Figure 6.) This clone contains a 1662 nucleotide cDNA insert encoding a 253 amino acid ORF and was cloned from a human T-cell lymphoma cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including B-cell lymphoma, corpus collosum, endometrial tumor, osteosarcoma, testes, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human tissues including spleen, lymph node, thymus, PBLs, heart, and a particularly strong signal in skeletal muscle and pancreas. A BLAST analysis of the amino acid sequence of HLTAH80 demonstrates that this clone exhibits approximately 35% identity and 55% similarity over the entire length of the A15 molecule.

1. 中心,一定指出的,如果的基本,也是是是这种的一个,但是是是这种的,是是是是是是是是是是是是是是是是是是是是是是是是是是是是是是是,也可以可以是是是是是是是

Since expression of A15 drops to undetectable levels when comparing immature T-cells to peripheral blood lymphocytes, it is thought that A15 may play a role in the development of T-cells. Furthermore, the MXS1(CCG-B7) gene which codes for A15 contains a number of triplet nucleotide repeats which have been associated with neuropsychiatric diseases such as Huntington's chorea, fragile X syndrome, and myotonic dystrophy. In addition, A15 appears to be expressed exclusively on T-cell acute lymphoblastic leukemia cell lines, including several derived from adult T-cell leukemia and those established by immortalization with human T-cell leukemia virus type 1 or Herpesvirus saimiri. Thus, clones HLTAH80 and/or HSBBF02 may also be involved in diseases caused by the expansion of repeats or chromosomal instability.

The cDNA clone HTPBA27 contains a 1345 nucleotide cDNA insert encoding a 238 amino acid ORF and was cloned from a human tumor pancreas cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including cerebellum, breast lymph node, osteosarcoma, adult testes, RS4;11 bone marrow cell line, microvascular endothelial cells, and others. A BLAST analysis of the amino acid sequence of HTPBA27 demonstrates that this

5

10

15

20

25

30

clone exhibits approximately 40% identity and 64% similarity with a glycoprotein termed CD53 over its entire length. CD53 is thought to be involved in thymopoiesis, since rat CD53 can be detected on immature CD4-8-thymocytes and the functionally mature single-positive subset, but cannot be detected on the intermediate CD4+8+ thymocytic subset of cells. The CD53 molecule has also been implicated as a component of signal transduction pathways in B cells, monocytes and granulocytes, rat macrophages, NK, and T cells. Moreover, as illustrated in Figure 7, HTPBA27 was recently confirmed as a TM4SF receptor. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997).) Calling the HTPBA27 polypeptide NAG-2, this group confirmed HTPBA27's status as a TM4SF receptor by showing that NAG-2 complexes with integrin and other TM4SF receptors. Thus, diseases caused by the failure of HTPBA27 to complex with integrin and other TM4SF receptors can be treated by administering HTPBA27. HTPBA27 can also be used to diagnose these diseases.

The cDNA clone HAIDQ59 contains cDNA insert encoding a 221 amino acid ORF that was cloned from a human epithelial cell induced with TNFa and INF cDNA library. The 5' end of HAIDQ59 is represented by the SEQ ID NO: 9, while the 3' end is represented by SEQ ID NO: 10. This clone appears in only two additional cDNA libraries in the HGS database. These two libraries were constructed from the human Jurkat T-cell line and human microvascular endothelial cells. A BLAST analysis of the amino acid sequence of HAIDQ59 demonstrates that this clone exhibits approximately 53% identity and 69% similarity over 226 amino acids of the CD9 TM4SF molecule. (See Figure 8.) It has been demonstrated that the CD9 molecule is involved in signal transduction pathways in platelets, as well as in cell adhesion in both platelets and pre-B-cell lines. Intriguingly, a monoclonal antibody (vpg15), which recognizes the feline homologue of CD9, has been shown to block infection by feline immunodeficiency virus (FIV). Furthermore, a recent study shows that cells expressing high levels of CD9 exhibited suppressed cell motility. Thus, HAIDQ59 may also be involved in signal transduction of blood cells.

The cDNA clone **HHFEK40** contains a 936 nucleotide cDNA insert encoding a 252 amino acid ORF and was cloned from a human fetal heart cDNA library. This clone appears once in the human fetal heart cDNA library and possibly in a hemangiopericytoma cDNA library. A BLAST analysis of the amino acid sequence of HHFEK40 demonstrated that this clone exhibits approximately 60% identity and 75% similarity over the entire length of a molecule designated PETA-3. (See Figure 9.) PETA-3 was originally identified as a novel human platelet surface glycoprotein termed gp27. Although PETA-3 is present in low abundance on the platelet surface, an anti-PETA-3 monoclonal antibody can stimulate platelet aggregation and mediator release.

5

10

15

20

25

30

Thus, HHFEK40 may function similar to PETA-3 to affect growth of blood cells. Administering polypeptides or fragments of HHFEK40 may be an effective treatment of blood disorders.

The cDNA clone HGBGV89 contains a 738 nucleotide cDNA insert encoding a 197 amino acid ORF and was cloned from a human gall bladder cDNA library. The only two additional appearances of this clone in the HGS database are in a normalized fetal liver cDNA library and in a fetal liver/spleen cDNA library. The cDNA clone HUVBB80 contains a 1071 nucleotide cDNA insert encoding a 201 amino acid ORF and was cloned from a human umbilical vein cDNA library. This clone appears in several additional cDNA libraries in the HGS database including prostate BPH, thyroid, and fetal liver/spleen. BLAST analyses of the amino acid sequences of HGBGV89 and HUVBB80 demonstrate that these clones exhibit approximately 49% identity and 65% similarity and 47% identity and 68% similarity, respectively, over the entire length of a molecule designated L6 surface protein or human tumor-associated antigen L6. (See Figures 10 & 11.) Moreover, another group has confirmed the TM4SF receptor homology of HGBGV89 by describing the protein as a putative transmembrane protein L6H. (See Genbank Accession No 2587054; see Figure 10.) The L6 cell surface antigen is highly expressed on lung, breast, colon, and ovarian carcinomas. Promising results of phase 1 clinical studies have been reported with an anti-L6 monoclonal antibody, or its humanized counterpart, suggesting that the L6 antigen may be an attractive target for monoclonal antibody-based cancer therapy.

日本の行の · 漢をおいて

In summary, there is a clear need for identifying and exploiting novel members of the TM4SF superfamily such as those described herein. Although structurally related, these factors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules, such as the novel potential members of the TM4SF superfamily detailed here, should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such factors may prove useful as therapeutics in an anti-tumor, diagnostic, or other capacity. Furthermore, factors such as the nine novel TM4SF superfamily-like molecules described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

Besides TM4SF receptors, receptors from other families are also described. For example, clone **HJACE54**, also called galectin 11, exhibits significant sequence identity to the rat galectin 5, the chicken galectin 3 gene, and the human galectin 8 genes. (See Figure 12.) The galectin 11 cDNA clone contains an 865 nucleotide insert which encodes a 133 amino acid ORF. The clone was obtained from a Jurkat T-cell G1 phase cDNA library. A BLAST analysis of the deduced amino acid sequence of HJACE54 demonstrates approximately 35% identity and 57% similarity to the amino

5

10

15

20

25

30

WO 98/31799 19 PCT/US98/00959

acid sequence of the rat galectin 5 gene. Expression of galectin 11 is quite limited in the HGS database. In fact, the only two additional ESTs in the HGS database which contain the HJACE54 sequence were found in human neutrophil and human infant adrenal gland cDNA libraries. Northern blot analyses have not been performed to examine expression patterns of the galectin 11 gene.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin 1 appears to inhibit cell adhesion of skeletal muscle presumably by galectin 1mediated disruption of laminin-integrin a₂b₁ interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bcl2, respectively. As a result, the discovery of a novel galectin, such as that encoded by HJACE54, is likely to be a valuable asset both diagnostically and therapeutically.

Additionally, a full-length nucleotide sequence of a novel human cDNA clone which encodes an apparent splice variant of the previously described human E48 antigen has recently been determined. (See Figure 13.) Clone HROAD63 contains a 441 nucleotide cDNA which encodes a 70 amino acid polypeptide. This novel clone exhibits significant sequence identity to several members of a relatively new family of cell-surface proteins termed the Ly6 superfamily. These members include murine and human SCA-2, rat Ly-6 (also termed ThB), and human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)]. The novel E48 splice variant was obtained from the HGS human stomach cDNA library. The clone is present in only a limited number of other HGS cDNA libraries including kidney cancer, keratinocyte, and tongue. An alignment of the nucleotide sequences of the human E48 and HROAD63 cDNAs demonstrates that the initial 168 and 178 nucleotides of E48 and HROAD63, respectively, are identical, with the exception of an additional 10 nucleotides of sequence at the extreme 5' end of the HROAD63 sequence. The

5

10

15

20

25

30

sequence of the two clones is also identical for an additional 229 nucleotides including the 3' end of the coding sequences and the entire 3' untranslated regions. The only divergence of nucleotide sequence in this region of the clones is the deletion of a single thymidine residue in the 3' UTR of the E48 cDNA. The major difference between the two nucleotide sequences is a 329 nucleotide deletion from the HROAD63 sequence. This deletion causes a shift in the HROAD63 reading frame and encompasses the translational stop signal used in the E48 clone. As a result, the carboxy terminal sequence of HROAD63 is radically altered with regard to that of E48 (as illustrated in Figure 13 by the obvious differences between amino acids 56-128 of E48 and 56-70 of HROAD63 in the amino acid alignment). The clinical presentation of disorders, including abnormal skin and hair phenotypes, may be attributed, at least in part, to a non-functional Ly6 superfamily member such as E48 or HROAD63. HROAD63 may also be involved in blood disorders, as seen with its homologues SCA-2 and CD59.

A novel prohibitin cDNA clone presented herein was originally identified in a human bone marrow cell line (RS4;11) cDNA library. The clone contains a 1066 nucleotide insert which encodes a 299 amino acid polypeptide. BLAST and BestFit analyses of the predicted amino acid sequence of HMWGS46 demonstrate a highly significant sequence identity to a murine protein termed IgM B-cell receptor associated protein (BAP)-37 (Genbank accession number X78683). The HMWGS46 amino acid sequence exhibits nearly perfect identity and similarity over the entire length of the murine BAP-37 sequence. (See Figure 14.) In addition, the full-length nucleotide sequences of HMWGS46 and BAP-37 exhibit at least 87% identical. The HMWGS46 clone also exhibits approximately 49% sequence identity and 85% sequence similarity to a human gene designated prohibitin. Finally, the HMWGS46 cDNA appears in a substantial number of HGS human cDNA libraries in addition to the bone marrow cell line cDNA library from which it was cloned. Some of the cDNA libraries in which this clone appears include keratinocytes, induced endothelial cells, activated neutrophils, synovial sarcoma, colon carcinoma cell line, Jurkat cell line membrane bound polysomes, epileptic frontal cortex, primary dendritic cells, and a number of others. The novel gene related to prohibitin and BAP-37 may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event. Thus, although the precise functional role of the prohibitin family members are less than clear, it is quite likely that such homologues are involved in such complex processes as development, senescence, and tumor suppression. Therefore a novel gene, such as HMWGS46, may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event.

A human cDNA clone encoding a novel epidermal growth factor receptor (EGFR)-like molecule is also disclosed. The novel EGFR-like cDNA clone presented herein was originally identified in an activated human neutrophil cDNA library. The

5

10

15

20

25

30

clone contains a 704 nucleotide insert which encodes a 168 amino acid polypeptide. A BLAST analysis of the predicted amino acid sequence of HNFGW06 demonstrates that this novel clone exhibits approximately 85% identity and 90% similarity to a protein designated epidermal growth factor receptor-related protein [Homo sapiens]. (See Figure 15.) The expression profile of the HNFGW06 clone in the HGS database indicates the existence of a fairly highly restricted expression pattern. In addition to the activated neutrophil library from which this clone was obtained, it also appears in the following HGS human cDNA libraries: synovial sarcoma, smooth muscle, placenta, and possibly primary dendritic cells.

PCT/US98/00959

The novel EGFR-like cDNA clone HNFGW06 may lead to a number of exciting possibilities for therapeutic and/or diagnostic treatments or reagents. For example, HNFGW06 may be involved in the onset of human breast cancers as well. In addition, due to the fact that TGF-a acts through binding to the EGFR, it is possible that HNFGW06 may also play a role in a variety of gastric processes including regulation of acid secretion, regulation of mucous cell growth, and protection against ethanol- and aspirin-induced injury to gastric tissues.

GENERATING POLYNUCLEOTIDES

Polynucleotides of the present invention encoding a receptor may be obtained using standard cloning and screening, from a cDNA library derived from mRNA in cells specified in Table 1 using the expressed sequence tag (EST) analysis (Adams, M.D., et al. Science (1991) 252:1651-1656; Adams, M.D. et al., Nature, (1992) 355:632-634; Adams, M.D., et al., Nature (1995) 377 Supp:3-174.) Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

The nucleotide sequence encoding a receptor polypeptide of SEQ ID NO:Y may be identical to the polynucleotide encoding SEQ ID NO:Y, or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:Y.

When the polynucleotides of the invention are used for the recombinant production of a receptor polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in

10

15

20

25

30

10

15

20

25

30

35

Gentz et al., Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Further preferred embodiments are polynucleotides encoding receptor variants comprising the amino acid sequence of receptor polypeptide of Table 1 (SEQ ID NO:Y) in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination.

The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 80%, and preferably at least 90%, and more preferably at least 95%, yet even more preferably 97-99% identity between the sequences.

Polynucleotides of the invention, which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:X or a fragment thereof, or to the cDNA insert in the plasmid deposited at the ATCC, or a fragment thereof, may be used as hybridization probes for cDNA and genomic DNA, to isolate full-length cDNAs and genomic clones encoding the receptor and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs) that have a high sequence similarity to the receptor gene. Such hybridization techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides. Preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides.

In one embodiment, to obtain a polynucleotide encoding the receptor polypeptide, including homologs and orthologs from other species, comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the SEQ ID NO:X or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to those of skill in the art. Stringent hybridization conditions are as defined above or, alternatively, conditions under overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

15

20

25

30

35

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease.

5 Vectors, Host Cells, Expression

The present invention also relates to vectors which comprise a polynucleotide or polynucleotides of the present invention, and host cells which are genetically engineered with vectors of the invention and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., BASIC METHODS IN MOLECULAR BIOLOGY (1986) and Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) such as calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, E. coli, Streptomyces and Bacillus subtilis cells; fungal cells, such as yeast cells and Aspergillus cells; insect cells such as Drosophila S2 and Spodoptera Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used. Such systems include, among others, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., MOLECULAR CLONING, A LABORATORY MANUAL (supra).

10

15

20

25

30

35

1000年,清晰的经验通过通常,增加强强调度等。 (1000年,1000年

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the desired polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If the receptor polypeptide is to be expressed for use in screening assays, generally, it is preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the receptor polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide; if produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

Receptor polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

Diagnostic Assays

This invention also relates to the use of receptor polynucleotides or polypeptides for use as diagnostic reagents. Detection of a mutated form of the receptor gene associated with a dysfunction will provide a diagnostic tool that can add to or define a diagnosis of a disease or susceptibility to a disease which results from under-expression, over-expression or altered expression of the receptor. Individuals carrying mutations in the receptor gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled receptor nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. (See, e.g., Myers et al., Science (1985) 230:1242.) Sequence changes at specific locations may also be revealed by

10

15

20

25

30

35

nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method. (See Cotton et al., Proc Natl Acad Sci USA (1985) 85: 4397-4401.) In another embodiment, an array of oligonucleotides probes comprising receptor nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability. (See for example: M.Chee et al., Science, Vol 274, pp 610-613 (1996).)

The diagnostic assays offer a process for diagnosing or determining a susceptibility to specific diseases through detection of mutation in the receptor gene by the methods described.

In addition, specific diseases can be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of receptor polypeptide or receptor mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit for a disease or susceptibility to a disease which comprises:

- (a) a receptor polynucleotide, preferably the nucleotide sequence of SEQ ID NO:X, or a fragment thereof;
 - (b) a nucleotide sequence complementary to that of (a);
- (c) a receptor polypeptide, preferably the polypeptide of SEQ ID NO:Y, or a fragment thereof; or
- (d) an antibody to a receptor polypeptide, preferably to the polypeptide of SEQ ID NO: Y.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Chromosome Assays

The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the

sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

10

15

20

5

Antibodies

The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them can also be used as immunogens to produce antibodies immunospecific for the receptor polypeptides. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against the receptor polypeptides can be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., Nature (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., Immunology Today (1983) 4:72) and the EBV-hybridoma technique (Cole et al., MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms including other mammals, may be used to express humanized antibodies.

30

25

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against receptor polypeptides may also be employed to treat diseases.

Vaccines

35

Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with a receptor polypeptide, or a fragment thereof, adequate to produce antibody and/or T cell immune response to protect said animal from a disease. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which

comprises, delivering a receptor polypeptide via a vector directing expression of the receptor polynucleotide in vivo in order to induce such an immunological response to produce antibody to protect said animal from diseases.

Further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a receptor polypeptide wherein the composition comprises a receptor polypeptide or receptor gene. The vaccine formulation may further comprise a suitable carrier. Since a receptor polypeptide may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations 10 suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation instonic with the blood of the recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, 15 sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The 20 dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

Screening Assays

5

The receptor polypeptide of the present invention may be employed in a screening process for compounds which bind the receptor and which activate (agonists) or inhibit activation of (antagonists) the receptor polypeptide of the present invention. Thus, polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See Coligan et al., Current Protocols in Immunology 1(2): Chapter 5 (1991).

The receptor polypeptides are responsible for many biological functions, including many pathologies. Accordingly, it is desirous to find compounds and drugs which stimulate the receptor on the one hand and which can inhibit the function of the receptor on the other hand. In general, agonists are employed for therapeutic and prophylactic purposes for such conditions and diseases. Antagonists may be employed for a variety of therapeutic and prophylactic purposes for such conditions and diseases.

In general, such screening procedures involve producing appropriate cells which express the receptor polypeptide of the present invention on the surface thereof.

25

30

10

15

20

25

30

Such cells include cells from mammals, yeast, Drosophila or E. coli. Cells expressing the receptor (or cell membrane containing the expressed receptor) are then contacted with a test compound to observe binding, or stimulation or inhibition of a functional response.

The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the receptor is detected by means of a label directly or indirectly associated with the candidate compound or in an assay involving competition with a labeled competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the receptor, using detection systems appropriate to the cells bearing the receptor at their surfaces. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed.

Further, the assays may simply comprise the steps of mixing a candidate compound with a solution containing a receptor polypeptide to form a mixture, measuring receptor activity in the mixture, and comparing the receptor activity of the mixture to a standard.

The receptor cDNA, protein and antibodies to the protein may also be used to configure assays for detecting the effect of added compounds on the production of receptor mRNA and protein in cells. For example, an ELISA may be constructed for measuring secreted or cell associated levels of receptor protein using monoclonal and polyclonal antibodies by standard methods known in the art, and this can be used to discover agents which may inhibit or enhance the production of the receptor (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues. Standard methods for conducting screening assays are well understood in the art.

Examples of potential receptor antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligand of the receptor, e.g., a fragment of the ligand, or small molecules which bind to the receptor but do not elicit a response, so that the activity of the receptor is prevented.

Thus in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for receptor polypeptides; or compounds which decrease or enhance the production of receptor, which comprises:

- (a) a receptor polypeptide, preferably that of SEQ ID NO:Y;
- (b) a recombinant cell expressing a receptor polypeptide, preferably that of SEQ 35 ID NO:Y:
 - (c) a cell membrane expressing a receptor polypeptide; preferably that of SEQ ID NO: Y; or
 - (d) antibody to a receptor polypeptide, preferably that of SEQ ID NO: Y.

SUBSTITUTE SHEET (RULE 26)

WO 98/31799 29 PCT/US98/00959

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Prophylactic and Therapeutic Methods

5

10

15

20

25

30

35

This invention provides methods of treating an abnormal conditions related to both an excess of and insufficient amounts of receptor activity.

If the activity of the receptor is in excess, several approaches are available. One approach comprises administering to a subject an inhibitor compound (antagonist) as described along with a pharmaceutically acceptable carrier in an amount effective to inhibit activation by blocking the binding of ligands to the receptor or by inhibiting a second signal, and thereby alleviating the abnormal condition.

In another approach, soluble forms of the receptor polypeptides still capable of binding the ligand in competition with endogenous receptor may be administered. Typical embodiments of such competitors comprise fragments of the receptor polypeptide.

In still another approach, expression of the gene encoding endogenous receptor can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered. (See, for example, O'Connor, J Neurochem (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Alternatively, oligonucleotides which form triple helices with the gene can be supplied. (See, for example, Lee et al., Nucleic Acids Res (1979) 6:3073; Cooney et al., Science (1988) 241:456; Dervan et al., Science (1991) 251:1360.) These oligomers can be administered per se or the relevant oligomers can be expressed in vivo.

For treating abnormal conditions related to an under-expression of the receptor and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates the receptor, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of the receptor by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells in vivo and expression of the polypeptide in vivo. For overview of gene therapy, see Chapter 20, Gene Therapy and

15

20

25

30

35

other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996).

5 Formulation and Administration

Peptides, such as the soluble form of receptor polypeptides, and agonists and antagonist peptides or small molecules, may be formulated in combination with a suitable pharmaceutical carrier. Such formulations comprise a therapeutically effective amount of the polypeptide or compound, and a pharmaceutically acceptable carrier or excipient. Such carriers include but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. Formulation should suit the mode of administration, and is well within the skill of the art. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds. Preferred forms of systemic administration of the pharmaceutical compositions include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if properly formulated in enteric or encapsulated formulations, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels and the like.

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of $0.1\text{-}100\,\mu\text{g/kg}$ of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide ex vivo, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

SUBSTITUTE SHEET (RULE 26)

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

																							<u> H</u>	M	A	\mathcal{L}	\mathcal{I}	<u>) </u>		
GCAG	TTO	CCT	SAG	AGA/	AGA	ACC	CTG	4GG/	AAC	AGA	GTI	rccc	TC	GCG	GCCC	TG	GCA	CCTC	CAA	ccc	CAC	AT/	TGC	TGC	TG	TGC	TGC	TGC	TG	20
CGTC	AAC	GGA	TC	CT	гст	TGG	GAC	TCC	TTG	TCT	GCA/	AGG	SAG	CGC	cecc	AC	CGT	GGAG	GTT	GGG	GT	TAT	FACG	ACC	AC	SAC	ACG	ACG	AC.	3 0
																							M	L	L	L	L	L	L	
cccc	TG	CTC	TGG	GG.	AGG	GAG	AGG	GTG	GAA	GGA	CAG	AAG	AGT	AAC	CGG	AGI	GAT	TAC	rcgo	TGA	ACG.	ATG	CAGA	GT	cc	GTG	ACCG	TGC	AA.	180
GGGG	AC	GAG.	ACC	CCC.	TCC	ctc	TCC	CAC	стт	CCT	GTC'	TTC	TCA	TTG	GCC1	rtc	CTA	ATG	AGC	GAC1	FGC	TAC	GTC1	TC A	AGG	CAC	rggc	ACC	TT	
P	L	L	W	G	R	ε	R	V	ε	G	٥	κ	s	N	R	K	D	Y	s	L	Т	M	0	S	s	٧	T	V	O	
GAGG	GC.	ATG	TGT	GTC	CAT	GTG	CGC	TGC	TCC	TTC	TCC	TAC	CCA	GTG	GAC	AGC	CAG	ACT	GAC	TCT	GAC	CCA	GTT	CAT	GGC	TAC	TGGT	TCC	GĠ	
CTC									_						_	_			_							_			_	270
Ε	G	M	С	٧	н	٧	R	С	s	F	s	Y	P	٧	D	s	0	T	D	s	D	P	٧	н	G	Y	W	F	R	
GCA	GG	AAT	GAT.	ATA	AGC	TGG	AAG	GCT	CCA	GTG	GCC.	ACA.	AAC	AAC	CCA	GÇT	TGG	GCA	GTG	CAG	GAG	GAA	ACT	CGG	GAC	CGA	ттс	AC	CTC	360
CGT	cc	TTA	CTA	TAT	TCG	ACC	TTC	CGA	GGI	CAC	CGG	TGT	TTG	TTG	GGT	CĠA	ACC	CGT	CAC	GTC	стс	CTT	TGA	GCC	CTG	GCT	AAG	STG	GAG	
A	G	N	D	ı	s	w	K	A	P	٧	A	T	N	N	P	A	W	A	٧	٥	Ε	Ε	T	R	Ð	R	F	н	L	
CTT	GG	GAC	CCA	CAG	ACC	:AAA	AAT	TGC	AC	CTG	AGC	ATC	AGA	GAT	GCC	AGA	ATG	AGT	GAT	GCG	GGG	AGA	TAC	ттс	TTT	CĢT	ATG	GAG	AAĄ	n=0
							_	+	_		_		+			-+-		TCA					_					•		450
	_	_	_					_			s	,	_			R		S	n		G		Y		F	R	м	Ε	κ	
L	G	U	P	0	,	K	N	L	•	_	-	٠	R	-				-			-				•	•				
				_		_							+			 -		CCT			_		-							540
CCT	TTA	TAT	TTT	ACC	TTA	ATA	TTT	ATA	CTO	GGTC	GAG	AGA	CAC	TTG	CAC	TGT	'AT(GGGA	GGA	GTC	TTO	AAC	TGA	CAC	TGA	\CAG	AAG	GTT	CET	
G	N	1	K	W	N	Y	K	Y	0	Q	L	s	٧	N	٧	T	Y	P	P	Q	N	L	T	٧	T	٧	F	٥	G	
GAA	GGC	ACA	GCA	TCC	AC/	AGCT	CTG	GGG	AAI	CAGC	TCA	TCT	CTI	TTCA	GTC	CTA	AGA(SGGC	CAG	TCT	сто	CGC	TTG	GTC	TG	rgçt	GTT	GAC	AGÇ	
											_		-			-	_	CCCG	_				_	_				•		630
								_	M									G					L				٧	_	s	
		•				A			14	S											_				Ī					
									_				+			- ÷		ACAG			_									720
TTA	GGG	GG/	ĊGG	TCC	GA	CTCG	ACC	TGG	AC	CTCC	TCA	GAC	TG	GGAC	CATG	GGG	SAG	TGTO	GGG	AGT	TTO	GG/	AGAC	CAT	GA	CCTC	GAC	GTT	CAC	
N	Ρ	P	A	R	L	s	W	T	W	R	S	L	T	L	Y	P	\$	Q	P	S	N	P	L	٧	L	Ε	L	Q	٧	
CAC	СТО	GGG	GAT	GAA	GG	GGAA	ATTO	ACC	:TG	TCGA	GCT	CAG	AAI	CTC	TCTG	GG	TTC	CCAG	CAC	GTT	TC	CTO	GAAC	CTO	TC	CCTO	CAA	CAG	GAG	810
GTG	GAC	ccc	CT/	CTI	CC	CCTI	AAG	TGG	AC	AGCT	CGA	GTC	TT	GAG	AGAC	cc	AAG	GGT	GTO	CAA	AG	GGA	CTTG	GAG	AG	GGA	GTT	GTC	стс	3.0
Н	L	G	0	Ε	G	Ε	F	Т	С	R	A	٥	N	s	L	G	S	0	н	٧	S	L	N	Ļ	S	L	a	۵	ε	

HMACR70 Page

				A T.C		CCT	GTA	TCA	GGA	стс	TTG	CTG	נננ	פרנ	GTCI	ຄຄຄ	GGA	ACT	GGA	GCC	ACA	GCC	CTG	GTC	TTC	CTC	TCC	TTC	rgt	
																							_				AGG	AAG	ACA	900
· · ·	т.	c	ĸ	м	ĸ	P	v	s	G	v	L	L	G	A	٧		G	Т	G	A	т	A	L	٧	F	L	s	F	C.	
•	,	G	^			•	•	•	Ī		_	_																		
			1								_		+			-+-		-			_		-			-		AAC	\rightarrow	990
CAG	TAG	AAG	TAA	CA1	CAC	TCC	AGG	ACG	TCC	TTC	TTT	AGC	CGT	TCT	GGT	CGT	CGC	CTG	CAC	ССТ	CTG	TAT	CCG	TAC	TTC	CTA	CGT	TIG	TGG	
٧	ı	F	1	٧	٧	R	s	С	R	ĸ	K	s	A	R	P	A	A	D	٧	G	D	I	G	M	K	0	A	N	T	
ATT	CAG	GGG	CTC	AGO	CTO	TCA	GGG	TAA	CTG	GAT	GAG	TCC	TGG	GCA	GAT	GAT	AAC	ccc	CGĄ	CAC	CAT	GGC	СTO	GCT	GCC	CĄC	TCC	CTC	AGG	1080
TAA	GTO	ccc	GAG	TC	GAC	SAGT	ccc	ATT	GAC	CTA	стс	AGG	ACC	CGT	CTA	CTA	TTG	GGG	GCT	GTO	GTA	CCC	GAC	CGA	CGC	GTG	AGG	GAG	TCC	1080
I	۵	G	Ļ	5	L	S	G	٠ (SA	ב כב	ED	NU	1: 1	ן ס																
GGA	GG#	AAG	AGA	GA	rcco	CAGT	ATG	CAC	ccç	TCA	GCT	TTC	ΑTΑ	AGG	GGG	AGC	CTC	AGG	ACC	TAI	rcc.	\GG1	CAA	AGA/	GC	CAÇC	AAC	TAA	GAG	1170
CCT	CC1	TTC	TCT	CT.	AGGO	STCA	TAC	GTO	GGG	AGT	CGA	AAG	TAT	TCC	ccc	TCG	GAG	TCC	TGG	ATA	AGGT	CCA	\GT1	rcti	CG	GTGG	TTO	STTA	стс	1170
TAC	TC	GAG	ATC	AA	GAT	ccc	AAG	TAA	GAA	AAT	GCA	GAG	GCT	CGG	GCT	TĢT	TTG	AGG	GTT	CAC	GA	ccc	TCC	CAGO	AA	AGG/	GT(TGA	GGC	1260
ATO	AG	сто	TAC	TT	CTA	GGGG	TTC	ATI	CTT	TTA	CGT	сто	CGA	GCC	CGA	AÇA	AAC	TCC	CAA	GTO	CTO	GGG	SAG	GTC	TT	TCC	CAC	SACT	ccc	1200
TGA	TT	CAG	TAC	AA	TTA	GCAG	scco	TCA	AATG	CTG	TGC	AAC	AAC	AC.	TCA	GAA	ACT	Γ ΑΤ Ί	rccj	CT.	TGT	TA	ACT	GAA	AAT	GC A	rgc	CTGA	TGA	1350
AC1	AA	GTC	ATO	TT	AAT	CGTO	GGG	AGI	TAC	GAC	ACC	TTG	110	TGI	AGT	CT	rga/	\TA/	\GG/	\GA/	ACA	GAT'	TGA	CTT	TTA	CGT	ACG(GACT	ACT	1350
-																														
c ¢ /	IAAI	TC1	cco	:11	TCC	CCAI	rcc#	ATO	GGT	CCA	CAC	TCC	:ccc	ccc	TGG	cçi	TCT(STAC	CCA	ACC.	ATT	CTC	cŢC	TGT	ACT	TCT	CTA	AGGA	TGA	1440
CC.	TT	GAGA	GGG	SAA	AGG	GGT	AGGT	TAC	SCC/	GGT	GTO	AGC	GGG	GGG	ACC	:GG/	4GA(CAT	GGG	T-G G	TAA	GAG	GAG	ACA	TGA	AGA	GAT	TCC1	TACT	1770
CT.	LCT	TTAC	AT	rcc	GAA	TAT	AGTO	AGA	ATTO	TAA	CGI	rga/	AAA	AAA	AAAA	AAA	444	444	۸		sec	ב כ	D	NO	:1)				
			_	_	+	ATA'		-											- 1.	497	-									

Page 1

	GGCACGAGTGGACAACCATCAGGGGAGCCAGGACACAGAGGGGGCAGAGCAAGTCAGCATTGGCGCCCCTTCCTCAGATCCTATCATCTT																													
CGG	AC	GAG	TGG	ACA	ACC	ATC	AGG	GAG	CCA	GGA	CAC	AGA	GGGI	GCA	GAG	CAA	GTC.	AGC	ATŢ	GGC	CCC	CT	CC.	TCA	GAT	ĊC.	TAT	CATO		
GCC	STG	CTC.	ACC	TGT	TGG	TAG	TCC	стс	GGT	ССТ	GTG	TCT	CCC	CGT	CTC	GTT	CAG	TCG	TAA	CCG	GGG	GA/	AGG.	AGT	CTA	SGG	ATAI	GTA		90
	-		; 			-+-		+	+				+						\rightarrow									ACT(+	180
																							м	A	Ε	I	н	T	Р	
TAT	TCT	TCC	TTG	AAG	AAA	CTG	TTA	TCT	TTA	CTC	AAT	GGC	TTC	GTG	GCT	GTG	TCT	GGC	AT:C	ATC	CTAC	3T T4	GGC	CTG	GGC.	ATT:	GGT	GGT	AAA	
			+ -						+		-+-		+			-+-		*	+					-				CCA.		270
Y	S	S	L	ĸ	ĸ	L	L	s	L	L	N	G	F	٧	A	v	s	G	ı	1	L	V	G	L	G	ı	G	G	ĸ	
	-+-		+			+		+			_	_	+			-+-		+					i 	$\overline{}$	_	 -		TGC		360
ACA	CCT	ccc	CGG	AGA	GAC	TGC	TTA	CAG	GAG	CCC	GAC	AGG	AGG	CGT	ATG	GAG	GAA	GTG	CAA	CCG	TTG	GAC.	ACG	GAC	CAC	TAC	CCT	ACG	TAG	
С	G	G	A	s	L	T	N	٧	Ł	G	L	s	s	A	Y	L	L	н	٧	G	N	L	С	L	٧	М	G	С	ı	
			÷			$\overline{}$						_	+			-+-		+	+							-		ATT	 +	450
TGC	CAT	GAC	GAA	CCG	ACA	CGG	ccc	ACC	ATA	CCT	CGC	TGA	TTT	стс	TCG	TCT	CCG	TGC	GAG	AAC	AAA	ACG	TAG	GAC	AGT	TAC	CAA	TAA	CAG	
T	٧	L	L.	G	С	A	G	W	Y	G	A	T	ĸ	E	S	R	G	T	L	L	F	С	ı	L	s	M	V	i	٧	
стс	ATC	ATG	GAA	GTT	ACA	GCT	GCC	ACÁ	GTG	GTC	CTT	CTT	TTC	TTT	CCA	ATT	GTT	GGA	GAT	GTG	GCC	TTG	GAA	CAC	ACC	TTC	GTG	ACC	CTG	
GAG	TAG	TAC	. 	CAA	TGT	CGA	CGG	TGT	CAC	CAG	GAA	GAA	AAG	AAA	GGT	TAA	CAA	CCT	CTA	CAC	CGG	AAC	t CTT	GTG	TGG	AAG	CAC	TGG	GAC	540
L	1	М	Ε	٧	T	A	A	7	٧	٧	L	L	F	F	P	I	٧	G	Ð	٧	A	L	Ε	н	T	F	٧	т	L	
											٠												c = 1		T00	T C T	~~~	CTC		
	-		+			-+-		+	\longrightarrow				+			-+-		+	+				+					GTG	_	630
TCC	TTC	TTA	ATG	TCT	CCA	ATG	TTG	CTC	GGT	CTG	CTG	ATA	AGA	TGT	GTC	ACC	TTG	AAC	CAG	TAC	CTC	TTC	GAT	TTC	ACG	ACA	CCC	CAC	TTA	•
R .	K	N	Y	R	G	Y	N	Ε	P	D,	D	Y	S	T	۵	W	N	L	٧	M	Ε	K	L	K	С	С	G	٧	N	
AAC	TAC	ACA	GAT	TTT	TCT	GGC	TCT	TCC	TTC	GAA	ATG	ACA	ACG	GGC	CAC	ACC	TAC	ccc	AGG	AGT	TGC	TGT	AAA	TCC	ATC	GĢA	AGT	GTG	TCC	700
TTG	ATG	TGT	CTA	AAA	AGA	cce	AGA	AGG	AAG	CTT	TAC	TGT	TGC	CCG	GTG	TGG	ATG	GGG	TCC	TCA	ACG	ACA	TTT	AGG	TAG	CCT	TCA	CAC	AGG	720
N	Y	T	D	F	s	G	s	s	F	Ε	M	т	T	G	н	T	Y	Р	R	s	С	c	ĸ	S	i	·G	s	٧	s	
TGT	GAC	GGA	CGC	GAT	GTG	тст	CCA	AAC	GTC	ATC	CAC	CAG	AAG	GGC	TGT	TTC	CAT	ΆΑΑ	.CTC	CTA	AAA	ATC	ACC	AAG	AC T	CAG	AGC	TTC	ACÇ	•
								+					+-						$\overline{}$				÷ ·					AAG		810
С	0	G	R	D	٧	S	P	N	٧	i	Н	Q	K	G	С	F	н	•		L		ı	7	ĸ	T		S		Т	
CTG	AGT	GGG	AGC	TCT	CTG	GGA	.GCT	GCA	GTG	ATA	CAG	AGG	ŢGG	GGG	тст	CGC	TAT	GTT	GCG	CAG	GCT	GGT	ÇTI	GAA	CTG	CŢG	GCC	TAA	AGC	PG TD 10:3
			- -			-+					-		+	\rightarrow		-+-	_			-,-	-		-					ATT		300
L	s	G	s	s	L	G	A	A	٧	i	a	R	W	G	s	R	Y	V	A	Q	A	G	L	Ε	L	L	A	٠ (

			_																					<u> []</u>	W	<u>'A</u>	صط	5		
GGC	STC	CCT	ÇTG	CCT	GCC	CĄC	TCA	GTG	GCĄ	ACA	CCC	GGG	AGC	TGT	TTT	GŢC	CTT	TGT	GGA	GCC	TCA	GCA	GTT	ccc	TCT	TTC	AGAA	CTC		90
CCG	CAG	GGA	GAC	GGA	CGG	GTG	AGT	CAC	CGT	TGT	GGG	ccc	TCG	ACA	AAA	CAG	GAA	ACA	CCT	CGG	AGT	CGT	CAA	GGG	AGA	AAG	TCTT	GA		90
																													•	
TGC	CAA	GAG	ccc	TGA	ACA	GGA	GCC	ACC	ATG	CAG	TGC	TTC	AGC	TTC	ATT	AAG	ACC	ATG	ATG	ATC	стс	TTC	AAT	TTG	CTC.	ATC:	TTTC	TG	TGT	
			: -	-		÷							+			—					 -					 -	AAAC		+	180
~~~	311		000	~~ .								,,,,,																		
									M	0	C	F	S	F	1	K	T	M	М	1	L	F	N	L	L	I	F	L	С	
GGT	GCA	GCC	CTG	TTG	GCA	GTG	GGC	ATC	TGG	GTG	TCA	ATC	GAT	GGG	GCA	TCC	: 111	CTG	AAG	ATC	TTC	GGG	CCA	CTG	TCG	TCC	AGT	3CC	ATG	
			<del></del>										+					<del>-</del>					<del>: -</del>	$\overline{}$		<del></del>	TCA		<del></del>	270
							_				_		_				_		<b>.</b>		_	_				_	_			
G	Α	A	L	L	A	٧	G	ı	W	٧	5	ı	D	G	А	5	1	L	K	ı	r	b	-	Ĺ	5	5	5	Α.	п	
CAG	TŢŢ	GTC	AAC	GTG	GGC	TAC	TTC	стс	ATC	GCA	GCC	GGC						CTT			CTG	GGC	ŢGC	TAT	GGT	GCT	AAG	AC T	GAG	360
GTC	AAA	CAG	TTG	CAC	CCG	ATG	AAG	GAG	TAG	CGT	CGG	CCG	CAA	CAC	CAG	AAA	ACG.	GAA	CCA	AAG	GAC	CCG	ACG	ATA	CCA	CGA	TTC	TGA	стс	500
n	E	v	N	v	c	Y	F	1	,	Δ	Δ	G	v	٧	v	F	A	L	G	F	L	G	С	Y	G	A	κ	т	Ε	
٠	•	•	••	•	•	•	•	_	•		•	Ī	·	·	•			_				-								
	-+-		<del></del>			÷		-			-+		+			<del></del> +-							÷	<del></del>		<del></del>	GTG	• • •	<del>+</del>	450
TÇG	TTC	ACA	CGG	GAG	CAC	TGC	AAG	AAG	AAG	TAG	GAG	GAG	GAG	TAC	SAAG	TAA	ACG/	ACTO	CAA	CGT	CGA	CGA	CAC	CAG	CGG	AAC	CAC	ATG	TGG	
s	K	С	A	L	٧	T	F	F	F	i	L	L	Ĺ	ī	F	ı	A	Ε	٧	A	A	A	٧	٧	A	L	٧	Y	T	
	4.70			C 4 C	TTC	CTC	٨٥٥	TTC	CTC	· C T A	CTC	CCT		· A T ſ	• ^ ^ C		AGA1	FT A T	reel	וזרר		CAA	CAC	TTC	`ACT		GTG	TGG	244	
			<del></del>			-+-		+			<del></del>		<del></del>			-		o					$\overline{}$			-+-	CAC	+	-+	540
161	IAL	LGA	L 1 L	616	AAG	GAL	166	AAC	GAL	LA I	LAL	GGA	1666	) i A (	3110	. 1 1 1	1617	4417	4007	1400	1010		CIG	,,,,,	3102			~~~		
T	.M	A	E	Н	F	L	T	L	L	٧	٧	Ρ	A	i	K	K	D	Y	G	S	Q	Ε	Đ	F	T	Q	٧	W	N	
ACC	ACC	ATG	AAA	GGG	стс	AAG	TGC	TGT	GGC	TTC	ACC	AAC	TAT	rac{	GGAT	ודדו	TGA	GGAC	CTCA	ACCC	TAC	TTC	AAA	GAG	SAAC	AGT	GCC	TTT	ĊCC	
	-+-		+			<del></del>		<del></del>					<del></del> -			-+-				<del></del>			+				CGG		$\overline{}$	630
										_				_	_	_	_	_	_	_		_		_				_	_	
T	Т	M	K	G	L	K	С	С	G	. F	T	N	Y	ī	D	F	E	D	5	۲	Ť	۲	K	£	N	3	A	r	r	
CCA	TTC	TGT	TGC	AAT	GAC	AĄC	GTC	ACC	AAC	ACA	GCC	AAT	GA/	ACC	TGC	AÇ	CAAI	GCA/	AAA	GCT	CAC	GAC	ÇAA	AAA	AGTA	GAG	GGT	TGC	TTC	720
GGT	AAG	ACA	ACG	TTA	CTG	TTG	CAG	TGG	TTG	TGT	CGG	TTA	CT1	TG	SACO	TG	STT	CGT	1110	CGA	GTO	CTO	GTI	TTT	CAI	cto	CCA	ACG	AAG	,20
D	_	_	r	M	n	N	v	Ŧ	N	т	Δ	N	£	т	_	т	ĸ	n	ĸ	Δ	н	۵	0	ĸ	٧	Ε	G	С	F	
•	•	·	٠	.,	U	,,	•	•	14	'	^	"	٠		٠	•	•	_		••	••	•	_		·	_	_	-		
AAT	CAG	CTT	TTG	TAT	GAC	ATC	CGA	ACT	AAT	GCA	GTC	ACC	GTO	GG	TGGT	FGT	GGC	AGC.	TGG	AATI	rgg(	GGG	:010	GAG	SCTO	GC1	GCC	ATG	TAA	810
TTA	GTC	GAA	AAC	ATA	CTG	TAG	GCT	TGA	TTA	CGT	CAG	TGG	CAC	CCA	ACCA	ACA	CCG	TCG	ACC1	TTA	CCC	:000	GAG	SC T(	CGAC	CGA	CGG	TAC	TTA	
N	Q	L	L	Y	0	1	R	T	N	A	٧	· T	٧	G	G	٧	A	, A	G	ī	G	G	L	Ε	L	A	A	M	N	
	_ 4									ــــــــــــــــــــــــــــــــــــــ														_		_				· SURI
ACA	CAG	GTA	CAT	AGA	CAT	GAC	GTT	AGA	TGT	TAT	TCA	GGT	GA/	AGA	CGG/	AGA	CGG	TGA'	TGA	CGAC	:GG	IGT/	ACC(	.11(	GAC	AC T T	CIC	CGT	GGG	
_					.,		_		-						۵		٥			,	P	н	G	N	c	F	F	Δ	P	

Päge 1

																								T	9	EF	ત્રા	٥		
<u>.</u>	AAA	AACA	AGG	TC	ccc.	ACAG	CAA	AGAA	AAA	GGA	ATA	GGA	TCA	AGA	GAT.	ACC	STG	CTG	CTG	GCAC	AG	CAA	GCA1	GA,	ATT	CGA	TGA	:TT	:A + 9	0
TTT	TT.	TTG	TCC	AGI	GGG	TGTC	GTT	TCT	III	сст	TAT	CCT	AG1	TCT	CTA	ŤG	CAC	CGAC	GAC	CGT	CTC	GTT	CGT	ACT	TAA	GCT	ACT	GAAG	ST	
																							1	4	N	_	M ·	T :	s ·	
							<b>_</b>						٠.			A T	CCT	GTGA	ררר	C A:C:	C A A	.TTA	TGT	ctc	ACC	TGC	מככ	TGT	ΑŤ	
GCAG	TT	CCG	GTGC	SCC.	AAT	TCTO	TGT	IGG	IGU	166	CAL	200	AL/	TTAC	CAA	H .	CCA	CACT	ccc	CTC	C T 1	ΓΔΑΊ	ACA	GAG	TG	ACC	GGG	ACA	<del>-+</del> TA	180
CGTC	AA	GGC	CAC	GG	TTA	AGA	CACA	ALC	ALL	.ALL	616	666	16	IIAL	LAA	. 1 A														
A	٧	•	٧			S				٧					_	Y		٧									٢	L	1	
CCAA	AC	AGC	CAG	ccg	CAA	GTC	CACC	TAG	TŢ	сте	GGA	ACÇ	CA	CCTA	GTT	ŢG	GTG	TCG	ATG	TGA	AT	GGGG	AGC	CTG	TG	CAG	AAAG	СТС	TG	270
GGT1	TG	TCG	GTC	GGC	GTT	CAG	GTGC	ATC	AAG	GGAC	ССТ	TGG	GT	GGAT	CAA	AC	CAC	AGC.	TAC	ACT	TA	ccc	STEG	GAC	AC	GTC'	TTTC	GAG	AC	_
														P				s					۵	P	٧	۵	K	A	L	
																	· A T C	GGC	TCC	:מרז	٠.	ΔΤΓ	ATGG	cer	1CG	GTT	CTCC	TAC	GG	
AAA	GAA	GGC	AAA	ACC	TTG	GGG	GCC/	1100	AG/	TAC	TACI	1116		CIGO	CAL	- TO	TAG	CCG	GAGO	CGA	GG	TAG	TACC	GC	rgc	CAA	GAG	ATO	+ :cc	360
																												v	G	
																		G										•	_	
GAA	TAC	CTO	TCT	ATI	TTC	TTC	TAC	GGAG	GÇ	TTT	ccc.	TCT	GG	GGA	GGC	770	STG	3777	ATC	ATT	TCA	GGA	TCT(	TC	TCC	GTG	GCA	GCA	GAA	450
CTT	ATO	GGAC	AGA	TA	AAG	TAAG	ATG	CCTC	CĠ	AAA	GGG	AAGA	ACC	CCT	CCG	AA(	CAC	CAAA	TAG	TAA	AGT	CCT	AGA	GAG	AGC	CAC	CGT	CGT	CTT	
Ε	Y	L	s	i	s	F	Y	G	G	F	P	F	W	G	G	L	W	F	ı	i	s	G	S	L	S	٧	A	A	E	
																		CAGT							GG	AGTO	ATA	стс	TTC	
AAI	CAI	500	- i - i	10	1 I A	4 A C G	CAC	CAC!	A G A	ccs	TCA	AAC	CCC	SAAC	TTG	TA	GCA	GTCA	CGT	TAG	ACC	AG/	CGT	CAA	CC.	TCAC	TAT	GAG	AAG	540
IIA	61																													
N	Q																	S												
ATC	AC	AGA	TCT	AAG	TAT	TCCC	CAC	CCA.	TAT	GCC	TAC	ccc	GA	CTAT	TAT	CC	TTA	cgc	TGG	GGT	GT(	SAA(	CCT	GGA	AT	GGC	SATI	101	GGC	630
TAC	TG	TCT.	AGA	TTC	ATA	AGG	GTG	GGT	ATA	CGG	ATG	GGG	CT	GATA	ATA	GG	AAT	GCG(	SACC	CCA	CA	CTT	GGGA	.001	TA	CCG	IAA	AGA		
i	Т	0	L	s	I	Ρ	Н	P	Y	A	Y	P	0	Y	Y	Ρ	Y	A	W	G	٧	N	Ρ	G	M	A	1	\$	G	
0.7.6		CCT	CCT		CTG	CCT	ר דו:	CAC.	T T 1	rego	ATC	GCA	TG	CGCA	ATC1	TC	CCA	CTT	rggc	TGC	CA	GTT	GGTC	TGC	TG	TCA	ATC	AGC	AAT	- 720
- G10	3L I	CCA	CC V	CAA	CAC	GGAI	GGAC	CTC	AA/	ACCO	TAG	CGT	AC	GCG1	AG/	AAG	GGT	GAA	ACCO	ACG	GT	CAA	CCAG	AC	AC	AGT	TAG	TCC	TTA	120
CAL	.uA	LGA	LLA	GAA	GAC	GUA												ł F									s			
V	L	. L	. V	F		. L	L																						<u></u>	
GT	GAG	TGT	CAT	CTA	TCC	AAA	CATO	TAT	GC	AGC/	AAC	CCA	GT	GAT	CAC	cc	AG	ACC	GGT	ACC	TC	ACC	ACC/	AAG	TTA	TTC	CAG	IGA	SAT(	810
CA	CTC	ACA	GTA	GAT	AGG	TTT	GTA	ATA	CG.	TCGT	TTT	GGT	CA	CTA	GTG	GGC	STCI	TOG	CCAG	CTG	GAG	TGG	TGGT	TTC.	AAT	AAG	GTC	ACT	LTA	•
v	9	5 V	, 1	١	/ F	, N	ı	Y	A	A	N	Ρ	٧	1	Т	f	- (	E P	٧	T	9	P	Р	S	١	' . S	S	Ε	i	

H ⁻	TPEF86	Page 2
CAAGCAAATAAGTAAGGCTACAGATTCTGGAAGCATCTTTCACTGGGACCAAAAGAAGTCCTCCCTTCCT		
GTTCGTTTATTCATTCCGATGTCTAAGACCTTCGTAGAAAGTGACCCTGGTTTTCTTCAGGAGGAGGGAAAGAG	CCGAAGGTATTGGGT	.C
OANK. (SEQIDNO: 21)		•
GTCGTTCCTGTTCTGACAGCTGAGGAAACGTCTCTCCCACTGTTTGTACTCTCACCTTCATTCTTCAATTCAG	TCTAGGAAACCATGCT	G + 990
CAGCAAGGACAAGACTGTCGACTCCTTTGCAGAGAGGGGTGACAAACATGAGAGTGGAAGTAAGAAGTTAAGTC	AGATCCTTTGGTACGA	vc
TTTCTCTATCAAGAAGAAGACAGAGATTTTAAACAGATGTTAACCAAGAGGGACTCCCTAGGGCACATGCATC	<del> </del>	_ 1ററേ
AAAGAGATAGTTCTTCTGTCTCTAAAATTTGTCTACAATTGGTTCTCCCTGAGGGATCCCGTGTACGTAG	TCGTGTATACACCCGT	TA
CCAGCCTCTGGGGCCTTGGCACACCCATTCGTGTGCTCTGCTGCATGTGAGCTTGTGGGTTAGAGGAACAAAT	ATCTAGACATTCAAT	:T → 1170
GGTCGGAGACCCCGGAACCGTGTGGGTAAGCACACGAGACGACGTACACTCGAACACCCAATCTCCTTGTTTA	TAGATCTGTAAGTTAG	SA 1170
TCACTCTTTCAATTGTGCATTCATTTAATAAATAGATACTGAGCATTCAAAAAAAA	SEQIDNO: 5	5)
AGTGAGAAAGTTAACACGTAAGTAAATTATTTATCTATGACTCGTAAGTTTTTTTT	-	

Päge 1

HSKRFO	7
CACGAGCAGGGTCTCGGGCTAGTCATGGCGTCCCCGTCTCGGAGACTGCAGACTAAACCAGTCATTACTTGTTTCAAGAGCGTTC	TGCTA 90
GTGCTCGTCCCAGAGCCCGATCAGTACCGCAGGGGCAGAGCCTCTGACGTCTGATTTGGTCAGTAATGAACAAAGTTCTCGCAAG	
MASPSRRLQTKPVITCFKSV	L L
ATCTACACTTTTATTTTCTGGATCACTGGCGTTATCCTTCTTGCAGTTGGCATTTGGGGCAAGGTGAGCCTGGAGAATTACTTTT	CTCTT 180
TAGATGTGAAAATAAAAGACCTAGTGACCGCAATAGGAAGAACGTCAACCGTAAACCCCGTTCCACTCGGACCTCTTAATGAAAA	GAGAA
I Y T F I F W I T G V I L L A V G I W G K V S L E N Y F	S L
TTAAATGAGAAGGCCACCAATGTCCCCTTCGTGCTCATTGCTACTGGTACCGTCATTATTCTTTTGGGCACCTTTGGTTGTTTTG	CTACC 270
AATTTACTCTTCCGGTGGTTACAGGGGAAGCACGAGTAACGATGACCATGGCAGTAATAAGAAAACCCGTGGAAACCAACAAAAC	GATGG
LNEKATNVPFVLIATGTVIILLGTFGCF	A T
TGCCGAGCTTCTGCATGGATGCTAAAACTGTATGCAATGTTTCTGACTCTCGTTTTTTTGGTCGAACTGGTCGCTGCCATCGTAC	
ACGGCTCGAAGACGTACCTACGATTTTGACATACGTTACAAAGACTGAGAGCAAAAAAACCAGCTTGACCAGCGACGGTAGCATC	CTAAA
CRASAWMLKLYAMFLTLVFLVELVAAIV	G F
GTTTTCAGACATGAGATTAAGAACAGCTTTAAGAATAATTATGAGAAGGCTTTGAAGCAGTATAACTCTACAGGAGATTATAGA	
CAAAAGTCTGTACTCTAATTCTTGTCGAAATTCTTATTAATACTCTTCCGAAACTTCGTCATATTGAGATGTCCTCTAATATCT	TEGGTA
V F R H E I K N S F K N N Y E K A L K Q Y N S T G O Y R	S H
GCAGTAGACAAGATCCAAAATACGTTGCATTGTTGTGGTGTCACCGATTATAGAGATTGGACAGATACTAATTATTACTCAGAA	
CGTCATCTGTTCTAGGTTTTATGCAACGTAACAACACCACAGTGGCTAATATCTCTAACCTGTCTATGATTAATAATGAGTCTT	TTTCCT
A V D K I Q N T L H C C G V T D Y R D W T D T N Y Y S E	K G
TTTCCTAAGAGTTGCTGTAAACTTGAAGATTGTACTCCACAGAGAGATGCAGACAAAGTAAACAATGAAGGTTGTTTTATAAAG	GTGATG
AAAGGATTCTCAACGACATTTGAACTTCTAACATGAGGTGTCTCTCTACGTCTGTTTCATTTGTTACTTCCAACAAAATATTTC	
F P K S C C K L E D C T P Q R D A D K V N N E G C F I K	v m
ACCATTATAGAGTCAGAAATGGGAGTCGTTGCAGGAATTTCCTTTGGAGTTGCTTGC	TACTGC 720
TGGTAATATCTCAGTCTTTACCCTCAGCAACGTCCTTAAAGGAAACCTCAACGAACG	
TILESEM G V V A G I S F G V A C F Q L I G I F L A	
CTCTCTCGTGCCATAACAAATAACCAGTATGAGATAGTGTAACCCAATGTATCTGTGGGCCTATTCCTCTCTCACCTTTAAGGAC	810
GAGAGAGCACGGTATTGTTTATTGGTCATACTCTATCACATTGGGTTACATAGAGACCCGGATAAGGAGAGATGGAAATTCCTC	( Anni •
LSRAITNNOYEIV. (SCG, ID NO: 22)	
A CONTRACTOR OF THE CONTRACTOR	STTGATT
CCAGGGGGGACACTTAATCTTTCAACGACCGACCTCTTGACTGTTGTGATGATGACTATCTGGTTTTTTGATGTGGTCATC	CAACTAA

																								1	ΓΑ	<u>H</u>	<u>30</u>			
CAC	SAG	CAT	TGC	GC.	TCT	CTC	GGT	SAGO	GC	AGC	ccc	CT	CTC	CGG	GCCG	GG	CCT	TCG	GGG	CCA	CC	GGC						GCA	TC	90
		GTA																												90
• • • •																														
																								п	6	u	С	L.	•	
ACC.	TCC	тсс	AAG	ACC	GTG	CTG	GTC.	TTT	тс	AAC	CTC	ATC	TTC	TGG	GGG	GÇA	GCT	GGC	ATŢ.	TTAT	rgc	TAT	STGG	GAC	SCC	TAT	GTCT	TC	ATC	180
		SAGG	1						-+	_				$\overline{}$		÷											-			100
								_					_		_							v	.,	_		v	,,	_	,	
T	S	S	K	T	٧	L	٧	F	L	N	L	i	F	W	G	A	A	G	ı	L	С	1	٧	G	A	1	٧	г	•	
ACT	TA ⁻	TGAT	GAC	TAT	GAC	CAC	TTC:	TTT	GAA	GAT	GTG	TAC	ACG	стс	ATC	CÇT	GCT	GTA	GTG	ATC	ATA	GCT	ĢTAC	GAC	GCC	CTG	CTT	TC	ATC	270
TGA	AT/	ACTA	CTG	ATA	CTG	GTG	AAG	AAA	CTT	CTA	CAC	ATG	TGC	GAG	TAG	GG.	CGA	CAT	CAC	TAG	TAT	CGA	CATO	СТС	ÇGG	GAC	GAA	AAG	TAG	210
														L				v	v	,		٨	v	c	Δ	,	,	F	ı	
Ţ	Y	_	0				F		_	_								•	•		٠	•	•	•		-	-	•	•	
ATT	GG	GCTA	ATT	GGC	TGC	TGT	GCC.	ACA	ATC	CGG	GAA	AGT	ÇGC	TGT	GGA	CTI	rgco	ACG	TTT	GTC.	ATC	ATC	CTG	CTC	TTG	GTT	TTT	GTC	ACA	360
TAA	CC	CGAT	TAA	CCG	ACG	AČA	CGG	TGT	TAG	GCC	CTT	TCA	GCC	SACA	ССТ	GA/	ACG	TGC	AAA	CAG	TAG	TAG	GAC	GAG	AAC	CAA	AAA	CAG	TGT	
	_		1	c	_	r	Δ	T	ı	R	Ε	s	R	c	c		Δ	τ	F	٧	I	ì	L	L	L	٧	F	٧	T	
1	6	_	-					-	-		_			-	-	_											_			
	٠.	TGT							_			_									_		_					_		450
CTT	CA	ACA	CAT	CAC	CAA	AAC	CCT	ATA	CAA	ATG	TCT	CGT	TT	CAC	CTT	TT	ACT	CAA	CTA	GCG	TCG	TAA	GTC	TTT	CAC	CATA	TTC	TGG	ATG	
F	٧	٧	٧	٧	٧	L	G	Y	٧	Y	R	A	K	٧	Ε	N	Ε	٧	O	R	s	1	۵	K	٧	Y	K	T	Y	
-	·																				701			c . c		- T A F	. T.C.A	C 4 C	TCC	
		AAC							_				_			-														540
TTA	CC	TTG	STTG	GGA	CTA	CGA	CGA	TCG	GCC	CGA	TAA	CTA	AT	ACAT	GTC	TC	TGT	CGAC	GTA	ACA	ACA	(CC)	IAA	616	111	JA I I	3AG 1	CIG	ALL	
N	G	Т	N	P	Đ	A	A	s	R	A	1	D	Y	٧	a	R	٥	L	н	С	C	G	ı	Н	N	Y	S	D	W	
								۸۲۲				ACT	ст	ccci	CTT	A C	CTC	rter	AGA	CAC	ΔC1	ופרו	. AGC	ΔΔΤ	TG	TAA'	recc	AGC	CTG	
																_					_				_					630
CTT	TT	ATG	TCTA	ACC	:AA	3111	CII	IGG	11	1116	i6 i C	. I C F	AL A	<b>,,,,</b> ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	46AA	· I C	GAL	GAL	3161		, 1 6/	4661	3166		IAC.	<b>~</b> ' ' '			GAC	
Ε	N	T	D.	W	F	K	ε	T	K	N	Q	S	٧	P	L	S	С	C	R	Ε	T	A	S	N	С	N	G	S	L	
000			***		·c Tí	- T A 1	rect	CAC	cci	STGI	rc a c	:cc1	r n T	AGT1	rete	:ΔΔ	GAA	GCT/	ACAA	AGAA	AT	CATO	SATG	CAT	GT	GAT	CTGG	GCC	GCA	
										-		_					_		_	Ī	_							•	CGT	720
CGG	iGī	GGG	AAGU	ille	i GAI	3A 1 A	ACGA			LALA	4C 1 C		NUA	ICA			C11	CGA			10	<b>.</b> , , , ,								
` <b>A</b>	Н	ı P	S	0	L	Y	A	Ε	G	C	Ε	A	L	٧	٧	K	K	L	a	Ε	i	M	M	Н	٧	1	W	A	A	
CTC		· A T T	TCC	rc 1	ΓΔΤ'	τιδι	ec TG	CTG	GGI	CATO	сте	TG	TGC	TTG	CATO	GT	GTT	GTG	CAG	AAGG	AG	TAG	AGAT	CCT	rgc	TTA	CGAC	CTO	TTC	
											${f}$					-			_		_							-	GAAG	810
GAC																														
			A	A		٥	1	1	C	м	- 1	C	Δ	C	1	٧	' L	C.	R	R	S	R	D	Ρ	Α	, Y	Ε	L	F	

	RAST.		
CACGAGCGCAGAGCTTGGGGCTTCCTTGGTCGCACCCACC	BAGCAC	CGCCTGGTC	90
GTGCTCGCGTCTCGAACCCCGAAGGAACCAGCGTGGGTGG			30
TCTTTCCTGTGGCCAGCCCAGAACTGAAGCGCTGCGGCATGGCGCGCGC	TGTTÇG	CCTTCAACC	
AGAAAGGACACCGGTCGGGTCTTGACTTCGCGACGCCGTACCGCGCGGACGGA	ACAAGC		
AGANAGGACAGGA			
MARACLUAVKTL	пг	AFN	
TGTTCTTCTGGCTGGGAGGCTGTGGCGTGCTGGGTGTCGGCATCTGGCCGCCACACAGGGGAGCTTCGCCAC	CGCTGT	сттсттсст	270
ACAAGAAGACCGACCCTCCGACACCGCACGACCCACAGCCGTAGACCGACC	GCGAĊA	IGAAGAAGGA	2.0
LFF W L G G C G V L G V G I W L A A T Q G S F A '	T - L	s s s	
TCCCGTCCCTGTCGGCTGCCAACCTGCTCATCATCACCGGCGCCTTTGTCATGGCCATCGGCTTCGTGGGCTGCC	TGGGTG	CCATCAAGG	
AGGGCAGGGACAGCCGACGGTTGGACGAGTAGTAGTGGCCGCGAAACAGTACCGGTAGCCGAAGCACCCGACGG			. 300
	ı 6	Δ 1 K	
FPSLSAANLLIITGAFV MAIGFVGC			
AGAACAAGTGCCTCCTGCTCACTTTCTTCCTGCTGCTGCTGCTGGTGTTCCTGCTGGAGGCCACCATCGCCATCC			
TCTTGTTCACGGAGGACGAGGGAAGAAGAAGGACGACGACGACCACAAGGACGAC	AGAAGA	AAGCGGATGT	•
ENKCLLLTFFLLLLÚVFLLEATIA!	L F	F A Y	
CGGACAAGATTGACAGGTATGCCCAGCAAGACCTGAAGAAAGGCTTGCACCTGTACGGCACGCAGGGCAACGTGG	GCCTC	ACCAACGCC	F 540
GCCTGTTCTAACTGTCCATACGGGTCGTTCTGGACTTCTTTCCGAACGTGGACATGCCGTGCGTCCCGTTGCACC	CCGGAG	TGGTTGCGG	
TO VIORYADODIKK GLHLY G T Q G N V	GL	T N A	
			_
GGAGCATCATCCAGACCGACTTCCGCTGCTGTGGCGTCTCCAACTACACTGACTG	<del></del>		, 000
CCTCGTAGTAGGTCTGGCTGAAGGCGACGACACCGCAGAGGTTGATGTGACTGAC	TGCGCC	CATGGACTG	Α .
W S I I Q T D F R C C G V S N Y T D W F E V Y N A	T R	V P D	
CCTGCTGCTTGGAGTTCAGTGAGAGCTGTGGGCTGCACGCCCCCGGCACCTGGTGGAAGGCGCCGTGCTACGAGA	ACGGTG	AAGGTGTGG	С
GGACGACGACCTCAAGTCACTCTCGACACCCGACGTGCGGGGGCCGTGGACCACCTTCCGCGGCACGATGCTCT			
GGACGACGTCAAGTCALTCTCGACACCCGACGTGCGGGGCCGTGGACCACCTTCCGCGGCACGACGACGACGACGACGACGACGACGACGACG	TOOUNG		_
S C C L E F S E S C G L H A P G T W W K A P C Y E	T V	K V W	
TICAGGAGAACCTGCTGGCTGTGGGCATCTTTGGGCTGTGCACGGCGCTGGTGCAGATCCTGGGCCTGACCTTCC	GCCATG	ACCATGTAC	T + 810
AAGTECTETTGGACGACCGGACACCCGTAGAAACCCGACACGTGCCGCGACCACGTCTAGGACCCGGACTGGAAGG			
. OF NILAVGIEG LCTALVOILG LTF			
			_
GCCAAGTGGTCAAGGCAGACACCTACTGCGCGTAGGCCGCCCACCGCCGGCTTCTCTGCCAAAAGGACGCCCACC			
CGGTTCACCAGTTCCGTCTGTGGATGACGCGCATCCGGCGGGTGGCGGCCGAAGAGACGGTTTTCCTGCGGGTG	CCCCTC	, I AUUGGEGT	li .
COVVKADTYCA. (SEQID'NO:24)			

·	45		TPRAZT	Page 2
CCACAGCTGCTTTTCCCACCACCAGCTTCGGTGTTCTGCCC	CATGCTGGGAGGAGGG	AGGGAGGGACAGGTGCC	TGGAGCCCCCGGAA	CC + 990
GGTGTCGACGAAAAGGGTGGTGGTCGAAGCCACAAGACGGG	GTACGACCCTCCTCCCT	CCCTCCCTGTCCACGG	ACCTCGGGGGCCTT	
CTGTTTCTGGAAGGCCCTAGCTCAGGTGGCTTCAGGGCCTC	CCGGACCCCCCCTGGGAC	GGGTGGCCACGTGCTG	GCTGCGGAACCCAG	GG 1080
GACAAAGACCTTCCGGGATCGAGTCCACCGAAGTCCCGGAG	GCCTGGGGGGGACCCTC	CCCACCGGTGCACGAC	CGACGCCTTGGGTC	
CAGGGGTGGGAGGGGCCTCCAGCACTTTTTATATTTACGTA	ATTCTCCAAAGCAGTGT	TCACACGGGAGCCAGCC	TGTGGCCCCCAGCT	TC 1170
GTCCCCACCCTCCCGGAGGTCGTGAAAAATATAAATGCAT	raagaggttTCGTCACA/	AGTGTGCCCTCGGTCGG	ACACCGGGGGTCGA	
CTGGAAAACAGGTTGGCGCTGGAGGAGCCGGGTCTTGGCAT	ICCTGGAGGTGGCCCCAI	CTGGTCCTGGTGCTCCA	GGCGGGGCCGTGGA	.cc 
GACCTTTTGTCCAACCGCGACCTCCTCGGCCCAGAACCGTA	AGGACCTCCACCGGGGT	GACCAGGACCACGAGGT	CCGCCCCGGCACCT	.cc
			(SEQ I	DIVO:8
CCTCACCTACATTCCATAGTGGGCCCGTGGGGCTCCTGGTG	CATCTTAATAAAGTGT	GAGCAGCAAAAAAAAA	MAAAAAAAAAA 13	- 345
GGAGTGGATGTAAGGTATCACCCGGGCACCCCGAGGACCAC	CGTAGAATTATTTCACA	CTCGTCGTTTTTTTTT	, , , , , , , , , , , , , , , , , , ,	•

																							t	<u> </u>	LC	$\mathcal{Q}$	59				
GCGC	:CG	CCG	GGC	CGC	AG	CAT	GGG	GC	GC	TTÇ	CGC	GGG	GGC	CTG	CGG	TGC	ATC.	AAG	TAC	CTĢ	CTG	CTT	GGC	TTC	۱AC	стĠ	CŢC	TTC	TGGC	TG	00
CGC	GC	GGC	CCG	GCC	TC	GTA	ccc	CG	CG	AAG	GCG	ccc	CCG	GAC	GCC.	ACG	TAG	TTC	ATG	GAC	GAC	GAA	CCG	AAG1	ГTG	GAC	GAG	AAG	ACCG		90
						м	G	:	R	F	R	G	G	L	R	С	t	K	Y	L	L	L	·G	F	N	L	L	F	W	۴.	
									•			_	_	-			•			_	_										
GCT	GGA	TCG	ĢCC	GTO	AT	TGC	TTT	TG	GA	CTA	TGG	TTT	CGG	TTC	GGA	GGT	GCC	ATA	AAG	GAĢ	TTA	TCA	TCA	GAG	GAC	AAG	TCC	CCA	GAGT	TAT	180
CGA	C T	AGC	CGG	CAC	STA	ACG	AAA	۱ÁC	СТ	GAT	ACC	AAA	GCC	AAG	CCT	CCA	CGG	TAT	TTC	стс	AAT	AGT	AGT	СТС	CTG	TTC	AGG	GGT	CTCA	ATA	
A	G	s	A	٧	I	A	F	:	G	L	W	F	R	F	G	G	A	i	K	Ε	L	s	S	E	D	K	s	<b>P</b>	Ε	Y	•
TTC	ΓΑΤ	GTG	GGG	CT	GTA	TGT	TCT	rgg	TT	GGA	GCC	GGG	GCC	ÇTG	ATG	ATG	GÇC	GTG	GGG	TTÇ	TTC	GGA	TGC	TGC	GGA	GCC	ATG	CGG	GAGT	rcĢ	270
AAG	ATA	CAC	CCC	GA	AT	ACA	AGA	VCC	AA	CCT	CGG	ССС	CGG	GAC	TAC	TAC	CGG	CAC	ccc	AAG	AAG	CCT	ACG	ACG	CCT	CGC	TAC	GCC	CTCA		270
F	Y	٧	G	L	Y	٧	Ł	-	٧	G	A	G	A	L	M	M	A	٧	G	F	F	G	С	c	G	A	M	R	E	S	
CAA	TGT	GTG	ÇTI	GG	ATC	ATŢ	TTI	ΓTΑ	cc	TGC	стс	CTG	GTG	ATA	TTT	GCT	GÇT	GAA	GTA	ACC	ACT	GGA	GTA	TTT	GCT	TTI	ATA	GGC	AAG	GG	360
GTT.	ACA	CAC	GAA	CC	TAG	TAA	AAA	AAT	GG	ACG	GAG	GAC	CAC	TAT	AAA	CGA	CGA	стт	CAT	TGG	TGA	CCI	CAT	ÀAA	CGA	AAA	TAT	CCG	TTC		300
a	С	٧	L	G	\$	F	F	•	T	С	L	L	٧	1	F	A	A	Ε	V	T	T	G	٧	F	A	F	ı	G	K	G	
GTA	GCT	ATC	CG/	(CA	TGT	TCĄ	GAC	CA	TG	TA]	GAA	GAG	GCT	ŢAC	AAT	GAT	TAC	CTT	AAA	GAÇ	AGG	GG/	AAA	GGC	AAI	GGG	AÇA	стс	ATC	ACC	450
CAT	CGA	TAC	GCI	GT	ACA	AGT	CTG	GT	AC	AT.	CTT	ctc	CGA	ATG	TTA	CTA	ATG	GAA	TTT	CTG	TCC	CCI	TTT	cce	TTA	cco	TĠT	GAG	TAG		750
٧	A	j	R	н	٧	a	1	Г	M	. <b>Y</b>	Ε	Ε	Ą	Y	N	D	Y	L	K	D	R	G	K	G.	N	G	T	Ł	i	T	
TTC	CAC	:TC/	ACA	LTT	TCA	GTG	CTO	STG	GA	AAA	GAA	AGC	TCC	GAA	CAG	GTC	CAA	CCT	TACA	TGC	CCA	AAC	GGAG	ÇTT	CT/	AGG/	ACAC	AAG	AAT	TGÇ	540
AAG	GTG	AG	TG	ΙAΑ	AGT	CAC	GAC	CAC	CT	TTI	CTT	TCG	AGG	CTI	GTC	CAG	GTT	GGA	TGT	ACG	GG1	TTO	CTC	GAA	GAT	rcc.	rgte	TTC	TTA		540
	υ		 T	-				-	c	ĸ	F	s	s	F	0	v	0	P	τ	c	ρ	ĸ	ε	L	L	G	н	ĸ	N	С	
r	п	3	•	r	_	•	Ì	•	•	``	-	Ū	J	_	•	•	•	•	•	Ū	•		-	_	Ī	Ī					
ATC	GAT	[GA/	AAT.	rga	GAC	CAT	AAI	TCA	GT	GT	AAG	CTC	CAG	CTO	ATT	GGA	ATT	GTO	CGGT	ATT	GG/	AAT.	TGCA	GGT	CTO	SAC	GATO	111	GGC	ATG	630
TAG	CT/	ACT	ΓŤΑ	ACT	C T G	GTA	TTA	AGT	CA	CA	TTC	GAG	GTC	ĠAC	TAA	CCT	TAA	CAC	SCCA	TAA	CCI	TA	ACGI	CCA	GAC	CTG	CTAC	AAA	CCG	TAC	
I	D	ε	I	Ε	T	1	i	i	s	٧	K	L	0	L	I	G	I	V	G	I	G	i	A	G	L	T	I	F	G	M	
ATA	TTC	CAG	ATI	GT	ССТ	CTG	сто	STO	CG	AT/	ACGA	AAC	TCA	CG/	AGAT	GTG	AŢA	TGA	AAGO	TAC	TTO	TA	CAT	AAA	AT.	TGC.	AAT	TAA	AGC	TTT	720
TAT	AAC	STC	STA	CA	GGA	GAC	GAC	CAC	GC	TAT	GCT	TTG	AGT	GC1	CTA	CAC	TAT	ACT	TTC	ATO	AAC	SAT	GTAC	:111	TA	ACG	TTA	TTAE	TCG		
ı	F	s	М	٧	L	. c	. (	5	A	i	R	N	s	R	D	٧	ı		(	C	Ì.		7	VO	, `	25	)	١			
_	-	GTT	+-		73	14 (	ĹS	Œ	Þ	J	.D	N	O`:	9	)												_				

Päge 1

HAT 0059	
AGTGTTTATGGGACTAAAAAACTTTTAACACCTTTTTAGGGGAAATATTTTGGTCCTATACAAAACATGTAAATATGCTTTATTACTTTC	90
AGTGTTTATGGGACTAAAAAACTTTTAACACCTTTTTAGGGGAAATATTTTGGTCCTATACAAAACATGTAAATATGCTTTATTACTTTC TCACAAATACCCTGATTTTTTGAAAATTGTGGAAAAATCCCCTTTATAAAACCAGGATATGTTTTGTACATTTATACGAAATAATGAAAG	
ATTTTCTGACCCTGCTGTAAACTACTGCAACCCTCACATCCCTCAAAGGGACTTTTATGTCAAACTCTTCTGTTTCTCCAAATATAAGGA TAAAAGACTGGGACGACATTTGATGACGTTGGGAGTGTAGGGAGTTTCCCTGAAAATACAGTTTGAGAAGACAAAGAGGTTTATATTCCT	180
AAAAAGACTAAAGCAAGAGATCTGGCAGTTGAAAATTGTGGGAAAGAGAATTTGTATGGGCACTGTATCTATGAAATACCTCATACTTAC TTTTTCTGATTTCGTTCTCTAGACCGTCAACTTTTAACACCCTTTCTCTTAAACATACCCGTGACATAGATACTTTATGGAGTATGAATG	270
GTTTACATGTTTTCCTAACTTTTTGTATTTTCTTGTATAGCCACCTAGAGAATTCTTCATAGATTAAGAACTACAGTTTTCACCACTTA	360
CAAATGTACAAAAGGATTGAAAAACATAAAAAGAACATATCGGTGGATCTCTTAAGAAGTATCTAATTCTTGATGTCAAAAGTGGTGAAT	
ACATAAGTAAAACAAAGTCCTTCATAATTTAACCATTAGCATCTTTGGCCAAACCAAAATAAAGAAAAGCATCTTCTCCTAGTTGTGTGTTTTGTATCATTTTGTTTCAGGAAGTATTAAATTGGTAATCGTAGAAACCGGTTTGGTTTTATTTCTTTTCGTAGAAGAGGATCAACACACAC	r - 450
TGTATTCATTTTGTTTCAGGAAGTATTAAATTGGTAATCGTAGAAACCGGTTTGGTTTTATTTCTTTTCGTAGAAGAGGATCAACACACAC	
GGGCAACAGAACAAGTTAAGGAAACAAAAATACTTATATACACAGAACAAAAATAATGTTCTTTTTATGCAAATCCCCTGTGAAAA	r + 540
CCCGTTGTCTTTGTTCAATICCTTTGTTTTTATGAATATATGTGTCTTGTTTTTATTACAAGAAAAATACGTTTAGGGGGACACTTTT	1
AAAATTTTCAATGTTTAAAAAAAAAAAAAAAAAAAAAAA	
TTTTAAAAGTTACAAATTTTTTTTTTTTTTTTTTTTTTT	

HHFEK40	_
TTCGGCACGAGCTGCGGGCGGCTGGGCGGCCCCGGGAGCCGCGCTCTCAGTCTCTCTAGGCGCAGTCCCTTCGCCGCTTCCGG	ļ
AAGCCGTGCTCGACGCCGCCACCCGCCGACCCGCGGGGCCCTCGGCGCGAGAGTCAGAGAGATCCGCGTCAGGGAAGCGGCGAAGGCC	
AGCCCCTGGCAGGGCCCAGAAGCCATGGCCCACTATAAGACTGAGCAGGACGACTGGCTGATCATCTACTTGAAGTATTTACTCTTTGTC	3O
TCGGGGACCGTCCCGGGTCTTCGGTACCGGGTGATATTCTGACTCGTCCTGCTGACCGACTAGTAGATGAACTTCATAAATGAGAAACAG	
MAHYKTEODOWLIIYLKYLLFV	
TTCAACTTCTTCTTCTGGGTCGGGGGGGGGCGCCGTCCTGGCTGTGGGCATCTGGACCCTGGTGGAGAAGAGTGGCTACCTCAGCGTCCTG	70
AAGTTGAAGAAGAAGACCCAGCCCCTCGTCGGCAGGACCGACACCCGTAGACCTGGGACCACCTCTTCTCACCGATGGAGTCGCAGGAC	
FNFFFW V G G A A V L A V G I W T L V E K S G Y L S V L	
GCCTCCAGCACCTTTGCCGCCTCCGCCTACATCCTCATCTTTGCGGGCGTACTTGTCATGGTGACCGGCTTCCTGGGCTTCGGTGCCATC	60
CGGAGGTCGTGGAAACGGCGGAGGCGGATGTAGGAGTAGAAACGCCCGCATGAACAGTACCACTGGCCGAAGGACCCGAAGCCACGGTAG	,
A S S T F A A S A Y I L I F A G V L V M V T G F L G F G A I	
CTCTGGGAGCGGAAGGGCTGCCTCTCCACGTATTTCTGCCTGTTGCTCGTCATCTTCCTGGTTGAGCTGGTGGCGGGAGTCCTGGCCCAT	
GAGACCCTCGCCTTCCCGACGGAGGGTGCATAAAGACGGACAACGAGCAGTAGAAGGACCAACTCGACCACCGCCCTCAGGACCGGGTA	50
GAGACCCTCGCCTTCCCGACGGAGGGGGCATAAAGACGGACAACGAGCAACGAGCAACGAGCAACGAGCAACGAGCAACGAGCAACGAGCAACGAGCAACGAGCAACGAGCAACGAGCAACGAGCAACGAGCAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACGAACAAC	
LWERKGCLSTYFCLLLVIFLVELVAGVLAH	
GTGTATTACCAGAGGCTGAGTGATGAACTGAAGCAGCACTTGAACCGGACTCTGGCTGAGAACTACGGGCAGCCGGAGCACGCAGATCAC	40
CACATAATGGTCTCCGACTCACTACTTGACTTCGTCGTGAACTTGGCCTGAGACCGACTCTTGATGCCCGTCGGCCTCGTGCGTCTAGTG	
V Y Y Q R L S D E L K Q H L N R T L A E N Y G Q P E H A D H	
GCCTCAGTGGACCGACTCCAGCAGGATTTCAAGTGCTGCGGAAGCAACAGCTCAGCCGACTGGCAGCACAGCACGTACATCCTGTTGCGG	
CGGAGTCACCTGGCTGAGGTCGTCCTAAAGTTCACGACGCCTTCGTTGTCGAGTCGGCTGACCGTCGTGTCGTGCATGTAGGACAACGCC	30
ASVORLOOFKEEGSNSSAUWUHSIIIE	
	20
CTCCGGCTCCCGGCGGTCCACGGCTGTCGACGACGTTCTGTCACCACCGCGCGACGCCGGTCGCCCGGGTGGGGAGGTTGTAGATATTC	
EAEGRQ V P D S C C K T V V A R C G Q R A H P S N I Y K	
GTGGAGGGAGGCTGCCTCACCAAGCTGGAGCAGTTCCTGGCCGACCACCTGCTGCTTATGGGGGCAGTGGGCATCGGGGTGGCCTGCCT	310
CACCTCCCTCCGACGGAGTGGTTCGACCTCGTCAAGGACCGGCTGGTGGACGACGACTACCCCCGTCACCCGTAGCCCCACCGGACGGA	110
VEGGCLTKLEOFLADHLLL MGAVGIGVACL	
	900
GTCTAGACGCCCTACCAAGAGTGGACGACGTGGTCTCCGAGGTCGCCGTAAAAATGATTACCGTTGGTGGAGGAGAAGGTTGACGG	
alcanvitacinariarh fy. (SCO.TD NO:26)	

GGAGTICTGTTGTACACCGTGTACGGTAGACGTTCC 936 SCO IDNO: 11

AGETTACTTTCACTCACCGCCTGTCCTTCCTGACACCTCACCATGTGTACGGGAAAAATGTGCCCGCTGTGTGGGGCTCTCCCTCATTACC TCGAATGAAAGTGAGGGGGACAGGAAGGACGTGGACGGGGGACACACCCCGGAGGGGACAAGGGCACACCCCGAGAGGGAAGGACGAGGAG	HGBGV89		
H C T G K C A R C V G L S L I T  CTCTGCCTGCTGCATTGTGGCCGAACGCCTCCTGGTACCTATGGGGAGAGCCTCCTGGACCAACCA	<del></del>		
CICTGGCCTCGTCTGCATTGTGGGCCAACGCCCTCCTGGTACCTAATGGGGAGACCTCCTGGACCAACCA	TCGAATGAAAGTGAGTGGCGGACAGGAAGGACTGTGGAGTGGTACACATGCCCTTTTACACGGGCGACACACCCCGAGAGGGAGTAA	rgg	
GAGACGGAGCAGACGTTACCACGGTTGCGGGAGGACCACCATGGATTACCCCTCTGGAGGACCTGGTTGTGGTTGGT	M C T G K C A R C V G L S L !	τ.	
GAGACGGAGCAGACGTTACCACGGTTGCGGGAGGACCACCATGGATTACCCCTCTGGAGGACCTGGTTGTGGTTGGT		<b>^</b> AA	
L C L V C I V A N A L L L V P N G E T S W T N T N H L S L O  CTCTGGCTCATGGGCGGCTTCATTGGCGGGGGCCTAATGGTACTGTGCTCCAGGGATTGCAGGCGTTCGGGCAAGGGCCGTCCCCGTTCCCGACGACA  Z70  CAGACCGAGTACCCGCCGAAGTAACCGCCCCCGGATTACCATGACACAGGTCCCTAACGTCGGCAAGCCCGTCCCCCGTTCCCGACGACA  V W L H G G F I G G G L H V L C P G I A A V R A G G K G C C  GGTGCTGGGTGCTGTGGAAACCGCTGCAGGATGCTGAGCCGAGGCCGTCCCCCGTTCCCGACGACA  GCAGACCCACGACACCTTTGGGCGACGTCCTACGACGCGAGCCCAGAAGAGAGGAGCCCCACGAACCACGGTAGATGACG  G A G C C G N R C R M L R S V F S S A F G V L G A I Y C L  GTGTCTGGACCTGGGCTCCGAAATGGACCCAGATGCTTAATGAACGGCGAGTGGGGCTACCACTTCGAAGACACCGCGGGAGCTTACTTG  CACAGACCTCGACCCGAGGCTTTACCTGGGTCTAAGGAATTACTTGCCGCTCACCCCGATGGTGAAGACTCTGTGGGGCCCCTCGAATGAAC  V S G A G L R N G P R C L M N G E W G Y H F E D T A G A Y L  CTCAAACCGCACTCTATGGGATCCCTTAGCGAGGGGGGGG		<del>+</del> 180	
CACGACCCACGACCCCTACGACGACTCATAGGACCCACGACGACCCCCCCGACGACCCCCCCC	GAGAL GGAGLAGALG TAALACESS TO COGGAGGALGALCATGGAT TACCCCTC TO GAGGACCTGG TTG GGT AGAT CGAAC	J	
CAGGACCGAGTACCCGCCGAAGTAACCGCCCCGGATTACCATGACACAGGTCCCTAACGTCGGCCAAGCCCGCCC	L C L V C I V A N A L L L V P N G E T S W T N T N H Ł S L	0	
V W L M G G F I G G G L M V L C P G I A A V R A G G K G C C  GGTGCTGGGTGCTGGAAACCGCTGCAGGATGCTGCGCTCGGTCTTCTCCTCGGGGTGCTTCGGGGTGCTTGGTGCCATCTACTG  CCACGACCCCAGGACACCTTTGGCGACGTCCTACGACGAGCCAAAGAGAGAG	GTCTGGCTCATGGGCGGCTTCATTGGCGGGGGCCTAATGGTACTGTGTCCAGGGATTGCAGCCGTTCGGGCAGGGGGCAAGGGCTGC	TGT 270	
GGTGCTGGGTGCTGTGGAAACCGCTGCAGGATGCTGCGCTCGGTCTTCTCCCTCGGCGTTCGGGGTGCTTGGTGCCATCTACTG CCACGACCCACGACACCTTTGGCGACGTCCTACGACGCGGAGCCAGAAGAGGGGGCCACAGCCCACGAACCACGGTAGATGACGG  G A G C C G N R C R M L R S V F S S A F G V L G A I Y C L  GTGTCTGGAGCTGGGCTCCGAAATGGACCCAGATGCTTAATGAACGGCGAGTGGGGGCTACCACTTCGAAGACCACCGGGGAGCTTACTTG CACAGACCTCGACCCGAGGCTTTACCTGGGTCTACGAATTACTTGCCGCTCACCCCGATGGTGAAGCTTCTGTGGGCGCCCTCGAATGAAC  V S G A G L R N G P R C L M N G E W G Y H F E D T A G A Y L  CTCAACCGCACTCTATGGGATCGGTGCGAGGCGCCCCCTCGCGTGGTCCCCTGGAATGTACTTCGCGTTGCTGGTGGCCGCCTCC GAGTTGGCGTGAGATACCCTAGCCACGCTCCCCGGGGGGAGCCACCACGGGGGACCTTACACTGCGAGAAAAAACACGGACCACCGGCGGAGG  L N R T L W D R C E A P P R V V P W N V T L F S L L V A A S  TGCCTGGAGATAGTACTGTGTGGGATCCAGCTGGTGAACGACCACTTGGTGTCTTCTGCGGCGATTGCAGGAAAAAACAGGACACCACCTTACACTGCGAGAAAAAACAGGACAACACCCTTACACTGCGAGAAAAAACAGGACAACACCCTTACACTGCGAGAAAAAACAGGACCACCCTTTTTTTT	CAGACCGAGTACCCGCCGAAGTAACCGCCCCGGATTACCATGACACAGGTCCCTAACGTCGGCAAGCCCGTTCCCGACG	ACA 270	
CCACGACCCACGACACCTTTGGCGACGTCCTACGACGCGAGCCAGAAGAGGAGCCGCAAGCCCCACGAACCACGGTAGATGACGE  G A G C C G N R C R M L R S V F S S A F G V L G A I Y C L  GTGTCTGGAGCTGGGCTCCGAAATGGACCCAGATGCTTAATGAACGGCGAGTGGGGCTACCACTTCGAAGACACCGCGGGAGCTTACTTG  CACAGACCTCGACCCGAGGCTTTACCTGGGTCTACGAATTACTTGCCGCTCACCCCGATGGTGAAGCTTCTGTGGCGCCCCTCGAATGAAC  V S G A G L R N G P R C L M N G E W G Y H F E D T A G A Y L  CTCAACCGCACTCTATGGGATCGGTGCGAGGCGCCCCCTCGCGTGGTCCCCTGGAATGTGACGCTCTTCTGCGCTGCTGGTGGCGCCCTCC  GAGTTGGCGTGAGATACCCTAGCCACGCTCCGCGGGGGAGCGACCACGGGGAACTTACACTGCGAGAAGAGCGACCACCGGCGGAGG  L N R T L W D R C E A P P R V V P W N V T L F S L L V A A S  TGCCTGGAGATAGTACTGTGGGGATCCAGCTGGTGAACGCGACCATTGGTGTCTTCTGCGGCGATTGCAGGAAAAAAACAGGACCACCCT  ACGGACCTCTATCATGACACACCCTAGGTCGACCACTTGCGCTGGTAACCACAGAAGAGCGCCGCTAACGTCCTTTTTTGTCCTGTGGGA  C L E I V L C G I Q L V N A T I G V F C G D C R K K Q D T P  CACTGAGGCTCCACTGACCGCCGGGTTACACCTGCCTCCTTCCT	V W L M G G F I G G G L M V L C P G I A A V R A G G K G C	С	
CCACGACCCACGACACCTTTGGCGACGTCCTACGACGCGAGCCAGAAGAGGAGCCGCAAGCCCCACGAACCACGGTAGATGACGE  G A G C C G N R C R M L R S V F S S A F G V L G A I Y C L  GTGTCTGGAGCTGGGCTCCGAAATGGACCCAGATGCTTAATGAACGGCGAGTGGGGCTACCACTTCGAAGACACCGCGGGAGCTTACTTG  CACAGACCTCGACCCGAGGCTTTACCTGGGTCTACGAATTACTTGCCGCTCACCCCGATGGTGAAGCTTCTGTGGCGCCCCTCGAATGAAC  V S G A G L R N G P R C L M N G E W G Y H F E D T A G A Y L  CTCAACCGCACTCTATGGGATCGGTGCGAGGCGCCCCCTCGCGTGGTCCCCTGGAATGTGACGCTCTTCTGCGCTGCTGGTGGCGCCCTCC  GAGTTGGCGTGAGATACCCTAGCCACGCTCCGCGGGGGAGCGACCACGGGGAACTTACACTGCGAGAAGAGCGACCACCGGCGGAGG  L N R T L W D R C E A P P R V V P W N V T L F S L L V A A S  TGCCTGGAGATAGTACTGTGGGGATCCAGCTGGTGAACGCGACCATTGGTGTCTTCTGCGGCGATTGCAGGAAAAAAACAGGACCACCCT  ACGGACCTCTATCATGACACACCCTAGGTCGACCACTTGCGCTGGTAACCACAGAAGAGCGCCGCTAACGTCCTTTTTTGTCCTGTGGGA  C L E I V L C G I Q L V N A T I G V F C G D C R K K Q D T P  CACTGAGGCTCCACTGACCGCCGGGTTACACCTGCCTCCTTCCT	***************************************		
G A G C C G N R C R M L R S V F S S A F G V L G A I Y C L  GTGTCTGGAGCTGGGCTCCGAAATGGACCCAGATGCTTAATGAACGGCGAGTGGGGCTACCACTTCGAAGACACCGGGGGAGCTTACTTG CACAGACCTCGACCCGAGGCTTTACCTGGGTCTACGAATTACTTGCCGCTCACCCCGATGGTGAAGCTTCTGGGCGCCCTCGAATGAAC  V S G A G L R N G P R C L M N G E W G Y H F E D T A G A Y L  CTCAACCGCACTCTATGGGATCCGTGGCGAGGCGCCCCCTCGCGTGGTCCCCTGGAATGTGACGTCTTCTCGCTGCTGGTGGCCGCCTCC  GAGTTGGCGTGAGATACCCTAGCCACCCTCCGCGGGGGAGCGCCCCCTCGGAATGTGACCTCTTCTCGCGAAAGACGCGACCACCCGGCGGAGG  L N R T L W D R C E A P P R V V P W N V T L F S L L V A A S  TGCCTGGAGATAGTACTGTGTGGGATCCAGCTGGTGAACGCGACCATTGGTGTCTTCTCGCGGCGATTGCAGGAAAAAACAGGACCACCCT  ACGGACCTCTATCATGACACACCCTAGGTCGACCACTTGCGCTGGTAACCACCAGAAGACGCGCGCTAACGTCCTTTTTTGTCCTGTTGGA  C L E I V L C G I Q L V N A T I G V F C G D C R K K Q D T P  CACTGAGGCTCCACTGACCGCCGGGGTTACACCTGCTCCTTCCT			
GTGTCTGGAGCTGGGCTCCGAAATGGACCCAGATGCTTAATGAACGGCGAGTGGGGCTACCACTTCGAAGACACCGCGGGAGCTTACTTG CACAGACCTCGACCCGAGGCTTTACCTGGGTCTACGAATTACTTGCCGCTCACCCCGATGGTGAAGCTTCTGTGGCGCCCCTCGAATGAAC  V S G A G L R N G P R C L M N G E W G Y H F E D T A G A Y L  CTCAACCGCACTCTATGGGATCGGTGCGAGGCGCCCCTCGCGTGGTCCCCTGGAATGTGACGCTCTTCTCGCTGCTGGTGGCCGCCTCC GAGTTGGCGTGAGATACCCTAGCCACGCTCCGCGGGGGGAGCGACCACCAGGGGACCTTACACTGCGAGAAGAGAGCGACCACCGGGGGAGG  L N R T L W D R C E A P P R V V P W N V T L F S L L V A A S  TGCCTGGAGATAGTACTGTGTGGGGATCCAGCTGGTGAACGCGACCATTGGTGTCTTCTGCGGCGGATTGCAGGAAAAAACAGGACACACCT ACGGACCTCTATCATGACACCCCTAGGTCGACCACTTGCGCTGGTAACCACAGAAGACGCGCGCTAACGTCCTTTTTTGTCCTGTTGGA  C L E I V L C G I Q L V N A T I G V F C G D C R K K Q D T P  CACTGAGGCTCCACTGACCGCCGGGGTTACACCTGCTCCTTCCT	CCACGACCCACGACACCIIIGGCGACGICCIACGACGCGAGCCAGAGAGGAGCCGCAAGCCCACGAACCACGGIAGAIGACGC		
CACAGACCTCGACCCGAGGCTTTACCTGGGTCTACGAATTACTTGCCGCTCACCCCGATGGTGAAGCTTCTGTGGCGCCCTCGAATGAAC  V S G A G L R N G P R C L M N G E W G Y H F E D T A G A Y L  CTCAACCGCACTCTATGGGATCGGTGCGAGGCGCCCCCTCGCGTGGTCCCCTGGAATGTGACGCTCTTCTCGCTGCTGGTGGCCGCCTCC  GAGTTGGCGTGAGATACCCTAGCCACGCTCCGCGGGGGAGCGACCACCAGGGGACCTTACACTGCGAGAAGAGAGCGACCACCGGCGGAGG  L N R T L W D R C E A P P R V V P W N V T L F S L L V A A S  TGCCTGGAGATAGTACTGTGTGGGATCCAGCTGGTGAACGCGACCATTGGTGTCTTCTGCGGCGATTGCAGGAAAAAAACAGGACACACCT  ACGGACCTCTATCATGACCACCCCTAGGTCGACCACTTGCGTGAACCACAGAAGACGCCGCTAACGTCCTTTTTTGTCCTGTGTGGA  C L E I V L C G I Q L V N A T I G V F C G D C R K K Q D T P  CACTGAGGCTCCACTGACCGCCGGGGTTACACCTGCTCCTTCCT	G A G C C G N R C R M L R S V F S S A F G V L G A I Y C L		
CACAGACCTCGACCCGAGGCTTTACCTGGGTCTACGAATTACTTGCCGCTCACCCCGATGGTGAAGCTTCTGTGGCGCCCTCGAATGAAC  V S G A G L R N G P R C L M N G E W G Y H F E D T A G A Y L  CTCAACCGCACTCTATGGGATCGGTGCGAGGCGCCCCCTCGCGTGGTCCCCTGGAATGTGACGCTCTTCTCGCTGCTGGTGGCCGCCTCC  GAGTTGGCGTGAGATACCCTAGCCACGCTCCGCGGGGGGAGCGACCACGGGGAACCACCTGCGAAAGAAGAGAGCGACCACCGGCGGAGG  L N R T L W D R C E A P P R V V P W N V T L F S L L V A A S  TGCCTGGAGATAGTACTGTGTGGGGATCCAGCTGGTGAACGCGACCATTGGTGTCTTCTGCGGCGAAAAAAAA	GTGTCTGGAGCTGGGCTCCGAAATGGACCCAGATGCTTAATGAACGGCGAGTGGGGCTACCACTTCGAAGACACCGCGGGAGCTTAC	TTG	
CTCAACCGCACTCTATGGGATCGGTGCGAGGCGCCCCCTCGCGTGGTCCCCTGGAATGTGACGCTCTTCTCGCTGCTGGTGGCCGCCTCC  GAGTTGGCGTGAGATACCCTAGCCACGCTCCGCGGGGGGAGCGACCACGGGGACCTTACACTGCGAGAAGAAGAGCGACCACCGGCGGAGG  L N R T L W D R C E A P P R V V P W N V T L F S L L V A A S  TGCCTGGAGATAGTACTGTGTGGGATCCAGCTGGTGAACGCGACCATTGGTGTCTTCTGCGGCGATTGCAGGAAAAAACAGGACACACCT  ACGGACCTCTATCATGACACACCCTAGGTCGACCACTTGCGCTGGTAACCACAGAAGACGCCGCTAACGTCCTTTTTTGTCCTGTGTGGA  C L E I V L C G I Q L V N A T I G V F C G D C R K K Q D T P  CACTGAGGCTCCACTGACCGCCGGGGTTACACCTGCTCCTTCCT	CACAGACCTCGACCCGAGGCTTTACCTGGGTCTACGAATTACTTGCCGCTCACCCCGATGGTGAAGCTTCTGTGGCGCCCTCGAATG		
GAGTTGGCGTGAGATACCCTAGCCACGCTCCGCGGGGGAGCGCCACCAGGGGACCTTACACTGCGAGAAAAAGAGGCGACCACCGGCGGAGG  L N R T L W D R C E A P P R V V P W N V T L F S L L V A A S  TGCCTGGAGATAGTACTGTGTGGGATCCAGCTGGTGAACGCGACCATTGGTGTCTTCTGCGGCGATTGCAGGAAAAAACAGGACCACCT ACGGACCTCTATCATGACACCACCCTAGGTCGACCACTTGCGCTGGTAACCACAGAAGACGCCGCTAACGTCCTTTTTTGTCCTGTGGGA  C L E I V L C G I Q L V N A T I G V F C G D C R K K Q D T P  CACTGAGGCTCCACTGACCGCCGGGTTACACCTGCTCCTTCCT	V S G A G L R N G P R C L M N G E W G Y H F E D T A G A Y	L	
GAGTTGGCGTGAGATACCCTAGCCACGCTCCGCGGGGGAGCGCCACCAGGGGACCTTACACTGCGAGAAAGAGCGACGACCACCGGCGGAGG  L N R T L W D R C E A P P R V V P W N V T L F S L L V A A S  TGCCTGGAGATAGTACTGTGTGGGATCCAGCTGGTGAACGCGACCATTGGTGTCTTCTGCGGCGATTGCAGGAAAAAACAGGACCACCT ACGGACCTCTATCATGACACCACCCTAGGTCGACCACTTGCGCTGGTAACCACAGAAGACGCCGCTAACGTCCTTTTTTGTCCTGTGGGA  C L E I V L C G I Q L V N A T I G V F C G D C R K K Q D T P  CACTGAGGCTCCACTGACCGCCGGGTTACACCTGCTCCTTCCT		TCC	
LNRTLWDRCEAPPRVVPWNVTLFSLLVAAS  TGCCTGGAGATAGTACTGTGTGGGATCCAGCTGGTGAACGCGACCATTGGTGTCTTCTGCGGCGATTGCAGGAAAAAACAGGACACACCT ACGGACCTCTATCATGACCACCACCTAGGTCGACCACTTGCGCTGGTAACCACAGAAGACGCCGCTAACGTCCTTTTTTGTCCTGTGGGA  CLEIVLCGIQLVNATIGVFCGCTGCTCCTTCCTGGACGCTCACTCCCTTGCTCGCTAGAATAAACTGCTTTGCGCTCTCA  TCACTGAGGCTCCACTGACCGCCGGGTTACACCTGCTCCTTCCT		+ 540	
ACGGACCTCTATCATGACACCCCTAGGTCGACCACTTGCGCTGGTAACCACAGAAGACGCCGCTAACGTCCTTTTTTGTCCTGTGGA  C L E I V L C G I Q L V N A T I G V F C G D C R K K Q D T P  CACTGAGGCTCCACTGACCGCCGGGTTACACCTGCTCCTTCCT	GAGTTGGCGTGAGATALLLTAGLLACGLTLLGLGGGGGGGCGCALLAGGGGGACLTTALACTGLGAGAAGAGGGCGALGACGACCACCGGCGG		
ACGGACCTCTATCATGACACCCCTAGGTCGACCACTTGCGCTGGTAACCACAGAAGACGCCGCTAACGTCCTTTTTTGTCCTGTGGA  C L E I V L C G I Q L V N A T I G V F C G D C R K K Q D T P  CACTGAGGCTCCACTGACCGCCGGGTTACACCTGCTCCTTCCT	LNRTLWDRCEAPPRVVPWNVTLFSLLVAA	S	
ACGGACCTCTATCATGACACCCCTAGGTCGACCACTTGCGCTGGTAACCACAGAAGACGCCGCTAACGTCCTTTTTTGTCCTGTGGA  C L E I V L C G I Q L V N A T I G V F C G D C R K K Q D T P  CACTGAGGCTCCACTGACCGCCGGGTTACACCTGCTCCTTCCT		CCT 630	
CACTGAGGCTCCACTGACCGCCGGGTTACACCTGCTCCTTCCT			
GTGACTCCGAGGTGACTGGCGGCCCAATGTGGACGAGGAAGGA	CLEIVLCGIQLV NATIGV F CGD C R K K Q D T	P	
GTGACTCCGAGGTGACTGGCGGCCCAATGTGGACGAGGAAGGA		TCA	
+ (SeQIDNO:27)		/20	
	`		
	+ (SeQIDNO:27)		
AAAAAAAAAAAAAAAC 738 (SCO IDNO: 12)	AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA		

Page
------

																							ł	<b>+</b> (	۷۷	Bi	186	0_		
GC	ACG	AGA	GAT	TGT	CGG	CTG	CGG	GTA	TAT	TCC.	AAT'	TCC	CG	TCT	CCTC	ATO	AAI	AŤG	AAG	TGA	AGG	GCT	CTG	ACC	CTG	GAA	GTG	GTT	CT	90
																												CAA		50
																													<u>.</u> .	
														_		_							_					ACA		180
TTC	GTC	CCG	TTT	TAC	ccc	AGA	GCC	TTC	ACA	CCT	CCG.	ACG	GAT	TCA	ACA	AACI	GAC'	TAA	SGC(	SAAG	CGT	SAAA	ACCI	CA	[AT]	AGC	:AC1	TTGT	AT	
				M	G	s	R	K	C	G	G	С	L	s	С	L	L	i .	Ρ	L	A	L	W	S	I	1	٧	N	ī	
ΓTΑ	TTO	TAT	TTC	CCG	AAT	GGG	CAA	ACT	TCC	TAT	GCA	TCC.	AGC	AAT	AAA	CŢC.	ACC.	AAC	TAC	STG	TGG	TAT:	TTT	SAAG	GGA/	ATC	TGT	TTCT	CA	270
AAT	AAC	ATA	AAG	GGC	TTA	ccc	GTT	TGA	AGG	ATA	CGT	AGG	TCG	TTA	TTT	GAG	TGG	TTG.	ATG	CAC	ACC	ATA	AAA	CTT	CCT	TAG		AAGA		270
		v	_		N	G	0	т	s	Y	Δ	S	S	N	K	L	T	N	Y	٧	W	Y	F	Ε	G	i	С	F	s	
_	L	1	r	r		_	·	•	٠													<b>.</b>	•••				*^^			
															_			_	_		_		<del>,</del>			-,-		AGC		360
CCC	TA	GTA	TAC	GAA	TAT	CAT	TGT	TGT	CAA	GAA	GAC	CAT	GAC	CTC	TTA	TTA	TTG	TTG	ATA	TTT	ACA	ACG	GTC	TCA	CII	HG	ACG	TCGT	111	
G	i	M	M	L	1	٧	T	T	٧	L	L	٧	L	Ε	N	N	N	N	Y	K	С	С	Q	S	Ε	N	С	S	K	•
44	\TA	TGT	GAC	ACTO	CTO	STCA	ATI	TATO	:771	тст	TCC	CTC	GGA	ATT	GCT	דיִד	TCT	GGA	TAÇ	TGC	CTG	GTC	ATC	TCT	GCC	TTG	GGT	CTT	GTC	450
				٠											_				_									GAA		450
r	Y		T		L	s	I	1	F	s	s	L	G	I	A	F	S	G	Y	c	L	٧	ı	s	A	L	G	L	٧	
•	•	-	-														ACT	CCT	CCV	CGT	TTC	CTT	ΔΩΔ	GAT	TCT	AGC	ΔΤΔ	TGG	ATT	
-																		_	_	_				_				ACC		540
GT	rcc	CGG	TAT	AAC	3GC(	GTG(	SGA/	AC I /	ACC	ALL		. A I A	1664	AAA,		CCG	1 1 1 1 1	CGA		56-										
0	G	P	Y	C	R	T	L	D	G	W	Ε	Y	A	F	Ε	G	T	A	G	R	F	L	T	0	S	S	i	W	į	
CA	GTG	CCT	GGA	ACC.	TGC	ACA'	TGT:	TGT	GGA	GTG	AAC	CATO	AT	TTT	1771	TCC	AT1	сто	ATA	ACC	CTC	AGT	GGG	CTI	CAA	GTG	ATC	ATC	TGC	·630
																												STAG		000
^			-	P	٨	Н	v	·v	E	w	N	1	ı	L	F	s		L		_		s		L	a	٧	I	i	С	
u			_	r	^		•	·	_																			`A.C.A		
																_							_					GACA		,
GA	GT#	GTC	TCA	TCA	GTA	CGT	TGA	TAG	GTT	CTA'	rga(	CAC	ACC.	TTC	GAT/	AAG	CAC	TAG	i A A 1	1611	,66,	166	IIAL	3 I A/				TGT		
L	1	R	٧	٧	М	Q	L	S	K	1	L	С	G	S	Y	S	٧	I	F	Q	P	G	I	ı	. (	_52	D	IJ	۱ (	ءُ:0،
																												TGCT		
AC	AAA	AGG	TAA	TAG	TTC	TGT	ACC	GGT	AGA	TAG	ATT.	TAT	AAT	ATA	GTT	GAC	ACA/	ATC1	rga/	CTO	ccc	STT	ATA	ACT.	TTT/	ACTA	4CC/	ACGA	AAG	
TO	CA	TTC	GTO	TTT	ATT	TGT	AAA	AAA	TTT	GCA	GTC	CTC	ACT	GCA	CAT	GC A	AGT	ATA	CAC	CC.	TTC	AT.	TTA	GTA.	TGT	<u> </u>	TA	AGTA	ATA	900
AC	GT/	AAAC	CAC	AAA	TAA	ACA	TTT	TTT	AAA	CGT	CAG	GAG	TGA	CGT	GTA	CGT	TCA	TAT	GGT	GG	AAG	GTA/	AATI	CAT.	ACA	AAA	AAT'	TCAT	TAT	•

IIIVERRO (°)

AGAAACCTTTATGCAATATGTATATTGCAACATTATTTAATATTCTGGAAAATTGGAAACACCCCAAAATTCTAACCTCAAA 1071
TCTTTGGAAATACGTTATACATATAACGTTGTAATAAATTATAAGACCTTTTAACCTTTGTGGGGTTTTAAGATTGAGTTT

Page 1

# HJACE54

ŢŢŢĢ	TC	GAG	ĢGC	AG	CAG	AGA	GT	ACC	CA	GCT	GGA	CA	TCC	TTC	CTO	GCT	GAT	GAG	cc	CCA	GGI	CTG	GAG	GTG	ccc	TGC	TC	AC	ATO	CT	CTT	ccc	90
AAAC	AC	CTC	ĊCG	TC	GTC	TCT	CA	TG	GGT	CGA	CCT	GT.	AGGA	AAAG	GA	CGA	CTA	CTC	GG	GGT	CC	GAC	CTC	CAC	GGG	ACG	AG	TG	TAC	GA	GAA	GGG	00
																	M	9	3	Р	R	L	Ε	٧	P	С	S	3	н	A	L	Ρ.	
CAGO	:G1	гстс	TCG	cc	TGG	GCA	GG	TC	ATC	ATA	GTA	CG	GGG	ACTO	GT	СТТ	GCA	AGA	\GC	CGA	AG	CAT	TTT	AC 1	FĢTO	SAGO	:c t	GA	ĢG(	GAC	CAG	GCŢ	
GTCC	:c/	AGAG	AGC	GG	ACC	CGT	cc	AG	TAG	TAT	CAT	GC	CCC.	TGAC	CA	GAA	CGT	TCI	rcs	GCT	TC	GTA	AAA	TG	CAC	TCC	GA	CT	CC	C T·G	GTC	CGA	180
۵	G	L	s	P	G	0	)	٧	ī	ŀ	. <b>V</b>	R	G	L	٧	L	. Q	. 8	Ξ :	Ρ	K	Н	F	T	٧	· ·S	L	-	R	D	a	A	
GCC	:A1	rgc1		GT	GAC	ACT	·CA	GGG	GCC	TCC	TTC	GC	AGA	CAGA	AAC	TCT	GGC	CTO	GGA	TCT	СĊ	CGC	TGG	GG	3ÇA(	GAAG	SA/	AAC	TG	ATC	TCA	GCĊ	070
			_		CTG		_								_		-		$\rightarrow$		-			_	<del>-</del> -	_	_		+				270
A	н	A	Р	٧	' Т	L		R	A	s	F	A	D	R	T	L	. А	١,	d	i	s	R	W	G	٥	ĸ	,	(	L	i	s	A	
CCC.	TTO	CCTO	<b>.TT</b>	ΓTΑ	ccc	CCA	\GA	\GA	TTC	TTT	GAG	GT	GCT	GCTO	CCT	GTI	CCA	GG	AGG	GAG	GĢ	CTG	AAG	CT	GĢC	GC T (	CA	ATG	GG	CAG	GGG	CTG	200
GGG	AA	GGA	AAA	AA1	GGG	GGT	CT	CT	AAG	AAA	CTO	CA	CGA	CGA	GGA	CAA	GGT	CC.	rcc	стс	cc	GAC	TTC	GA	ccg	CGA	GT.	TAC	ĊС	GTC	ccc	GAC	360
P	F	L	F	١	P	C	1	R	F	F	Ε	٧	L	L	Ļ	F	. 0	1	Ε	G	G	L	ĸ	L	A	L	ļ	N	G	٥	G	L	
ccc	GCI	CAC	AGO	: A ]	GAA	CCA	١GC	AG	GCC	сте	GAG	CA	GCT	GCG(	GGA	GC1	rccg	GA'	TCA	GTG	GA	AGT	GTO	:CA	GCT	CTA	CT	GTG	TC	CAC	TCC	TGA	
			1		CTT		_		+					<del></del>		+	o						<del></del>		-+-		٠-		+-		AGG	ACT	450
G	A	T	s	ı	1 N		3	a	A	L	Ε	0	L	R	ε		_ R	≀	ı	s	G	s	٧	a	L	Y	1	С	٧	н	s	<u>.</u>	DID No:2
					SAAA						- ۸ ۸ (	: A C	TCA	מרר.	A C T	רנו	- r A G	cc	רכו	·cac	:тс	TCC	TCC	cc	TCA	TTA.	AA	CCA	TC	CAC	CTG	AAC	
					TTT				+					-+-		<del></del>				—							-		+-		-		540
	'-	<b>.</b>																															
ACC	AG	CAC	AŢC.	AG	GCC	TG	311	TCA	CCI	сто	GGG	TC	ACG	AGA	CTG	AG	CTA	\CA	GG	AGC	TT	rgg	CCI	TGA	GGG	AAG	GC	AC/	AAG	AGT	GCA	AAG	630
TGG	TC	GTG	TAG	TC	CGG	ACC	CA/	AGT	GG/	AGAC	ccc	AG	TGC	TCT	GAC	TC	AGAT	rgt	CC.	rcg/	<b>AA</b>	ACC	GG	<b>ACT</b>	cċc	TTC							
																			_														
			- : -		TGCA		_		-									_			_	_			_	-	_		Ţ				120
CAA	GG	AGC	TTG	AG	ACGT	GGA	AA(	GGA	GG1	rGGT	rcc:	CG	GAC	CCT	ATA	(CCI	SAGE	ilA	GA	-66/	AAL	31 C	.461	3AL	616	ALG	I'G	AG	161	<b>L</b> 1 <b>L</b>		IICA	
GTT	GT	AGA	CTA	AC.	AAAG	ATA	AC1	tcc	AAA	AAT	ACA/	\TG	GCT	TĄA.	AGA	ĄTI	GTGG	STC	AT'	TTA'	TTO	TT	TAT'	TAT	TTA	TTT	ΑT	110	G Ţ G	GTO	AAA	TAA	810
CAA	CA	TCT	GAT	TG	TTTC	TAT	TG/	AGG	TT	TA:	rg T	AC	CGA	ATT	тст	TAI	CACC	AG	TA	AAT	AAC	SAA	TA	ATA	AAT	AAA	TA	AA	CAC	CAC	STTI		
																														_			
ATA	AA	TAA	GGT	TΑ	TTT/	TT	TA/	AAA	AAA	AAA	AAA	AAA	AAA	AAA	AAA	AA.	AAA	AAA	AA	A - 80	65	(5	æ	C	IC	N	C	):	14	ł )	•		
TAT	TT	ATT	rr.	ΔΤ	ΔΔΔ1		AT.	TTT	TTI	TTT	TTT'	<b>1</b> 11	TTT	TTT	TTT	TT	TTT	TTT	TT	T			_	•	_ ,	•							

HROAD63

Page 1

GCACGAGAGACGACATCAGAGATGAGGACAGCATTGCTGCTCCTTGCAGCCCTGGCTGTGGCTACAGGGCCAGCCCTTACCCTGCGCTGC 90
CGTGCTCTCTGCTGTAGTCTCTACTCCTGTCGTAACGACGAGGAACGTCGGGACCGACACCGATGTCCCGGTCGGGAATGGGACGCGACG

M R T A L L L A A L A V A T G P A L T L R C

CACGTGTGCACCAGCTCCAGCAACTGCAAGCATTCTGTGGTCTGCCCGGCCAGCTCTCGCTTCTGCAAGACCACGAACACAGTGGAGCCT
GTGCACACGTGGTCGAGGTCGTTGACGTTCGTAAGACACCAGACCGGCCCGGTCGAGAGCGAAGACGTTCTGGTGCTTGTTGTCACCTCGGA

H V C T S S S N C K H S V V C P A S S R F C K T T N T V E P

CTGAGGGCTTCCCCGAAAGTCTGGGACCAGGTCCAGGTGGGCATGGAATGCTGATGACTTGGAGCAGGCCCCACAGACCCCACAGAGGAT 270
GACTCCCGAAGGGGCTTTCAGACCCTGGTCCAGGTCCACCCGTACCTTACGACTACTGAACCTCGTCCGGGGTGTCTGGGGTGTCTCCTA

LRASPKVWDQVQVGHEC. (SCQIDNO: 30)

GAAGCCACCCCACAGAGGATGCAGCCCCCAGCTGCATGGAAGGTGGAGGACAGAAGCCCTGTGGATCCCCGGATTTCACACTCCTTCTGT
CTTCGGTGGGGTGTCTCCTACGTCGGGGGTCGACGTACCTTCCACCTCCTGTCTTCGGGACACCCTAGGGGCCTAAAGTGTGAGGAAGACA

HMW6546

AGC	GG	GC	C C I	GAA	CCC	TCG	TG1	GAA	GG	GTO	GC A	GT.	ACC	TAA	GCC	:GG	AGC	GG	GGT	AG	AGG	CGG	GC	CGG	SC A	CCC		TC	GA	CCT	CCA	GTG	GEC	90
TCG	CC	CG	GGI	CTT	GGG	AGC	:ACA	ACTI	CC	CAC	CGT	CA	TGG.	ATT	CGG	SCC.	TCG	CC	CCA	ATC	TCC	GCC	CG	GCC	:G1	GGC	SGGA	AAG	AC T	GGA	GGT	CAC	GGC	,
	٠.										-			_	+		+			_	+		_		•					_	_	-	AAG	180
GGC	CG	GA	GT	TCT	AGT	CTO	STAC	CGC	GT	CT.	TG/	AC	TTC	CTG	AAC	CG	CCC	TG	CCC	SAC	GGC	CGG	CC	CG(	GGG	CCC	CCG	TAC	CCG	TGC	CGG	GAC	TTC	
							M	A	0	) (	N	L	K	D	L	A	. (	3	R	L	Ρ.	A	G	1	P	R	G	M	G	T	A	L	K	
CTG	Tī	GC	TG	GGG	GCC	GGC	GCC	CGT	GC	CT.	AÇ	GT	GTG	CGC	GA/	ATC	TG	rg1	TC	ACC	GTO	GA/	GG	CGI	GGC	AC	AGA	GCC	ATC	TTC	TTC	AAT	rcee	270
GAC	AA	CG	AC	ccc	CGG	CCC	3CG(	SCAC	CG	GA	TĠ	CA	CAC	GCG	ĊT.	TAG	AC/	AC/	AAG	TGG	CAC	CT	rcc	GC	CCC	TG	TCT	CGG	TAG	AAG	AAG	TTA	AGCC	
L	L		L	G	A	G	A	٧	A		Y	G	٧	R	Ε	S	. 1	٧	F	T	٧	ε	G	i (	G	H	R	A	Ì	F	F	N	R	
ATC	GG	STG	GA	GTG	CAG	CAC	GGA	CAC	ΓΑΤ	CC	TĢ	GCC	GAG	GGC	ĊT.	TCA	CT.	TC/	AGG.	ATC	CCI	TG	STT	CC.	AG	TAC	ÇCC.	ATT	ATC	TAT	GAC	AT	TCGG	360
TAG	CC	AC	CT	CAC	GTO	:GT(	ccT	GTG/	AT A	GG.	ACC	GG	стс	CCG	GA.	AGT	GA/	AG	rcc	TAG	GG/	ACC	CAA	GG	TC	ATG	ĠGG	TAA	TAC	ATA	CTG	TA	AGCC	
ı	6	3	G	٧	۵	۵	D	T	1		L	A	Ε	G	L	н	. 1	F	R	1	P	W	F		Q	Y	P	Į	I	Y	٥	ı	R	
GCC	AC	GAC	:cT	ÇGA	AAA.	ATI	СТС	стсі	ccc	TA	CA	GGC	TCC	AAA	GA	CCT	AC	AG/	A T.G	GTG	AA.	TAT	CTC	cc	TG	GA	GTG	TTG	TC	rcg/	ccc	AA.	TGCT	450
CGG	T	TG	GA	GCT	TT	ГТАІ	GAG	GAG	GGG	AT	GT	CCG	AGG	TTI	ĊTI	GGA	TG	TC.	TAC	CAC	ŤT.	ATA	GAG	GG	AC1	SCT	CAC	AAC	AGA	AGC1	GGG	TT	ACGA	
A	ı	R	P	R	K	I	s	s	F		T	G	s	K	0	L		Q	M	٧	N	t	S	6	L	R	٧	L	S	R	P	N	A	
CAG	G/	AGC	:11	ÇC1	AG	CAT	GTA	CCA	GCG	cc	TĄ	GGG	CTG	GAC	TA	CGA	GG	AA	CGA	GTO	TT	GCC	GTC	CA	TT	GTC	AAC	GAG	GT	SCTO	AAC	SAG	TGTG	540
GTC	C.	TCC	AA	ĠG/	TC	GTA	CAT	GGT	CGC	GG	ΑŤ	ccc	GAC	CTO	AT	GCT	CC	T T	GĊT	CAC	:AA	CGG	ÇAG	GT	AA	CAG	TTG	CTC	CA	CGA	STT	CTC	ACAC	
Q	(	Ε	L	Ρ	s	M	Y	a	F	2	L	G	L	D	Y	Ε	•	Ε	R	٧	L	P	5	8	Ī	٧	N	ε	٧	L	K	S	٧	
GTO	G	cc/	AAG	TTO	:AA	TGC	стс	ACA	GC 1	ΓGA	TC.	ACC	CAC	CGG	gc	CCA	GG	TA	TÇC	ст	STT	GAT	cco	cc	GG	GAG	CTG	AC/	GA	GAG	GGC	CAA	GGAC	: - 630
CAC	:CI	GGT	гтс	AAC	TT.	ACG	GAG	TGT	CG/	ACT	AG	TGG	GTC	GC	:cG	GGT	CC	AT.	AGG	GA	CAA	CTA	ĠGC	CGG	CC	CTC	GAC	TGT	CT	CTC	CCG	GTT	CCTG	1
٧		A	K	F	N	A	S	Q	L	-	ı	T	a	R	A		1	٧	S	L	L	I	F	R	R	Ε	L	T	E	R	A	K	D	
TTO	A	GCI	CTC	ATO	:cT	GGA	TGA	TGT	GGC	CA	TÇ.	ACA	GAG	CTI	ĢĀG	CTI	ΙTΑ	GC	CGA	GAG	STA	CAC	AGO	CTG	СТ	GTA	GAA	GCC	:AA	ACA	AGT	GGC	CCAG	- 720
AAG	37	CG	GAC	TA	GA	CCT	ACT	ACA	ccc	GGT	AĞ	TGT	СТО	GA	TC	GA/	AAT	CG	GĊT	CT	CAT	GTG	TC	GAC	GA	CAT	ĊTI	rcgo	377	TGT	TCA	CCG	GGTC	:
F		S	L	1	L	D	D	V	,	4	i	T	ε	L	s		•	S	R	Ε	Y	T		Å	A	٧	Ε	A	K	۵	٧	A	۵ ،	
CA	36	AGI	GCO	CAI	SCG	GGC	CCA	ATT	CTI	TGG	TA	GAA	AAA	AGC:	۸Ą۸	.GC	\GG	AA	CĄG	CG	GCA	GAA	ΑA	TTC	TG	CAG	GCC	GAG	GGG	TGA	GGC	CGA	GGC	r - 810
GT	cc	TC	CGO	GT	GC	ccg	GGT	TAA	GA/	ACC	ΑŤ	CTI	TTI	CG	rtt	CG1	rcc	TT	GTC	GC	CGT	CTT	TT	AAC	AC	GTO	ccc	GC T	ccc	ACT	CCG	GCT	CCG	1
٥		Ε	A	Q	R	A	. a	F	į	L	٧.	Ε	K	A	K	. (	נ	Ε	Q	R	Q	K		ı	٧	a	A	Ε	G	Ε	A	E	<b>.</b> A	
GCI	CA	AG	ΑTO	3C T	TGG	AGA	AGC	ACT	GAG	GCA	AG	AAC	:cc:	rgg	CTA	CAI	ГСА	AA	CTI	CG	CAA	GAT	TC	GAG	SCA	GC	ÇAC	GAA'	TAT	CTC	CAA	GAC	GAT	. 000
																																	CTA	
					_	_					v	N	٥	c	٧	,	ı	ĸ	ı	Q	ĸ	i	,	R	A	A	Q	N	1	S	K	1	r i	

HMWF546

Page 2

GCCACATCACAGAATCGTATCTATCTCACAGCTGACAACCTTGTGCTGAACCTACAGGATGAAAGTTTCACCAGGGGAAGTGACAGCCTC
CGGTGTAGTGTCTTAGCATAGATAGAGTGTCGACTGTTGGAACACGACTTGGATGTCCTACTTTCAAAGTGGTCCCCTTCACTGTCGGAG

A T S Q N R I Y L T A D N L V L N L Q D E S F T R G S D S L

ATCAAGGGTAAGAAATGAGCCTAGTCACCAAGAACTCCACCCCCACAAGAAGTGGATCTGCTTCTCCAGTTTTTGA
TAGTTCCCATTCTTTACTCGGATCAGTGGTTCTTGAGGTGGGGGTGTTCTTCACCTAGACGAAGAGGGTCAAAAAACT

IKGKK (SOOJDNO: 3)

HIPF. WOL

GGCA	CGA	GAT	GAC	ATC	ACT	AAG	TGG	CCG	ATC	ŢGC	ACA	GAG	CAG	GCC	AGG	AGC.	AAC	CÁC	AÇA	GGC	TTC	CTG	CAC	ATG	GAC	TGC	GAG	ATC	AA	90
CCGT	GCT	CTA	CTG	TAG	TGA	T.T.C	ACC	GGC	TAG	ACG	TGT	стс	GTC	CGG	TCC	TCG	TTG	GTG	TĠT	CCG	AAG	GAĊ	GTG	TAC	CTG	ACG	CTC	TAG	ΤŤ	30
																		•						H	D	С	Ε	ı	ĸ	
GGGC	CGC	ccç	TGC	TGC	ATC	GGC	ACC	AAG	GGC	AGC	TGT	GAG	ATC	ACC	ACC	Ċee	GAA	TAC	TGT	GAG	TTC	ATG	CAC	GGC	TAT	TTC	CAT	GAG	GĄ	180
ccc	GCG	GGG	ACG	ACG	TAG	CCG	TGG	TTC	CCC	TCG	ACA	CTC	TAG	TGG	TGG	ccc	CTT	ATG	ACA	CTC	AAG	TAC	GTG	CCG	ATA	AAG	GTA	CTC	СŤ	100
G	R	P	С	С	· 1	G	T	K	G	s	C _.	ε	ī	T	T	R	Ε	Y	С	ε	F	М	н	G	Y	, <b>F</b>	Н	Ε	Ε	
AGCA	ACA	CTC	TGC	TCC	CAG	GTG	AGG	CGA	GGC	AGG	CCT	GGĄ	GTA	GTG	GAG	GAG	AGG	ACG	CTG	GGC	ATG	GCA	GCC	TGC	TGG	ĢGC	CGG	GGC	TC	270
TCGT	TGT	GAG	ACG	AGG	GTC	CAC	TCC	GCT	ccc	тсс	GGA	cct	CAT	CAC	CTC	СТС	TCC	TGC	GAC	CCG	TAC	CGT	cgg	ACG	ACC	CCG	GCC	CCG	AG	270
A	T	L	С	s	Q	٧	R	R	G	R	P	G	V	٧	Ε	Ε	R	T	L	G	M	A	A	С	W	G	R	G	s	
ACG						+			-+-		+	$\rightarrow$				+					<del></del>			<del></del>		+ -			<del>-+</del>	360
TGC	TGA	GGG	AGG	GTA	CAG	CCT	CGG	AGT	СТО	SAGT	CCG	ACG	AAG	ACC	CCG	CGA	CTC	GTG	GTA	TAC	GGG	TAA	GGG	TCC	CACG	TGA	CAA	AAC	CT	
R	. <b>T</b>	P	s	Н	٧	G	A	S	D	\$	G	С	F	W	G	A	Ε	Н	Н	M	P	1	P	R	С	T	٧	L	D	
CAA	GTG	TGT	TGG	GCT	GCT	GCC	TTC	CTC	AAC	ССТ	GAG	GTC	CCA	GAT	CAG	TTT	TAC	AGG	TCT	GGC	TGT	CTC	TTT	TCC	TAC	ATG	TTG	GG1	AA	450
GTT	CAC	ACA	ACC	CGA	CGA	ĊGG	AAG	GAG	TŤ	GGA	CTC	CAG	GGT	CTA	GTC	AAA	ATG	TCC	AĞA	CCG	ACA	GAG	SAAA	AGC	SATO	TAC	AAC	CCA		
K	٧	С	W	A	A	Ά	F	Ü	N	P	Ε	٧	P	0	a	F	Y	R	s	G	С	L	F	s	Y	M	L	G	K	
GAG	STCC	TCA	ATG	ccc	CCG	AAC	CCC	ACC	cçı	GTG	ATC	GAC	ACC	CAG	GCG	GAC	ccc	TGG	GGA	AAG	GTI	CCI	rggg	CCA	AGGG	ŢAT	GGT	CGG	TÇ	540
CTC	AGG	AGT	TAC	GGG	GGC	TTG	GGC	TGG	GG/	CAC	TAC	CTG	TGG	GTC	CGC	CTG	GGG	ACC	CC1	TTC	CAA	GG/	CCC	GGT	rcco	ÀΤΑ	CC/	GCC	AG	340
R	s	s	М	Ρ	P	N	P	τ	P	٧	M	D	Ţ	٥	A	D	P	W	G	K	٧	Ρ	G	P	G	Y	G	R	s	
CAA	сто	ccq	AAG	ACT	ACT	GCT	CCI	GAA	GŢ	STCT	GGA	TGA	AGG	CCG	CTG	ÇCT	GGI	GTG	TCC	CTC	ccc	CAC	TGT	GGG	STG	AC1	GCC	CTO	GĢ	630
GTT	GGAC	GGC	TTC	TGA	TGA	CGA	GGA	CTI	CAC	AGA	CCT	ACT	TCC	GGC	GAC	GGA	CC	CAC	AGC	GAG	GGG	GTO	ACA	ccc	CAC	TGA	CGC	GA	scc	030
N	L	Р	K	T	Ŧ	A	Ρ	ε	٧	s	G	. '	B	3	IC	1	<b>W</b> .	3	2)	)		•								
TGT	:TT0	TGG	GTC	TTT	TCA	AAT	GAC	ATO	יסם:	rgaa	GGG	GAC	сто	GAG	GAA	GGT	'GG1	rcce	GCT	GGC	ACC	CT	ATCO	E .	704	\S	9	I!	וכ	NO:T
ACA	SAAC	ACC	CAG	AAA	AGT	TTA	CTO	TAC	GG	CTT	ccc	CTG	GAC	CTC	CTT	CCA	CCA	GGC	CGA	CCC	TG	GA'	TAGO		, 57					

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

and the language of the mineral community of the second community of the secon	ad to in the description
A. The indications made below relate to the microorganism referr on page 13, line	Table 1
011 prog 2	Further deposits are identified on an additional sheet
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an auditional sheet
Name of depositary institution	•
American Type Culture Collection	
and an	-1
Address of depositary institution (including postal code and count	· · · · · · · · · · · · · · · · · · ·
12301 Parklawn Drive Rockville, Maryland 20852	·
United States of America	·
	L
Date of deposit May 16, 1997	Accession Number 209053
C. ADDITIONAL INDICATIONS (leave blank if not applicable	e) This information is continued on an additional sheet
·	
D. DESIGNATED STATES FOR WHICH INDICATIONS A	RE MADE (if the indications are not for all designated States)
	·
,	·
E. SEPARATE FURNISHING OF INDICATIONS (leave blan	nk if not applicable)
	Bureau later (specify the general nature of the indications e.g., "Accession
The indications listed below will be submitted to the international in Number of Deposit")	Suicau later (specify the general range c by the Daniel Co. Co.,
For receiving Office use only	For International Bureau use only
This sheet was received with the international application	This sheet was received by the International Bureau on:
	-
Standard Sta	Authorized officer
Authorized officer Yvette Sirum	
IAPD-PCT Operation:	
WAR OF 8762	

Form PCT/RO/134 (July 1992)

REC'D

2 3 FEB 1998

#### INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

	WIPO PCT
A. The indications made below relate to the microorganism referm on page13, lineTable	ed to in the description able 1 .
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution	
American Type Culture Collection	
Address of depositary institution (including postal code and country 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	(ער
Date of deposit	Accession Number
May 16, 1997	209053
C. ADDITIONAL INDICATIONS (leave blank if not applicable	e) This information is continued on an additional sheet
of the deposited microorganism will be mathemention of the grant of the European application has been refused of withdraw the issue of such a sample to an expert a sample (Rule 28(4) EPC).  D. DESIGNATED STATES FOR WHICH INDICATIONS A	patent or until the date on which the n or is deemed to be withdrawn, only by nominated by the person requesting the RE MADE (if the indications are not for all designated States)
E. SEPARATE FURNISHING OF INDICATIONS (leave blank	
Number of Deposit")	Bureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
This sheet was received with the international application	This sheet was received by the International Bureau on:
Authorized officer versa simple	Authorized officer
Particulal Specialist Particular Operations	
MARCHANI OCC.	

Form PCT/RO/134 (July 1992)

#### INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism reference on page 13 ,/in/ Tab	7
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution  American Type Culture Collection	
Address of depositary institution (including postal code and count 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	(ירי)
Date of deposit May 16, 1997	Accession Number 209054
C. ADDITIONAL INDICATIONS (leave blank if not applicable	This information is continued on an additional sheet
D. DESIGNATED STATES FOR WHICH INDICATIONS A	RE MADE (if the indications are not for all designated States)
E. SEPARATE FURNISHING OF INDICATIONS (leave bla	nk if not applicable)
	Bureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
This sheet was received with the international application	This sheet was received by the International Bureau on:
Authorized officer Vietto Status  Parallocal Specialist  APD-PCT Operations	Authorized officer

### 59/2

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

REC'D	2 3 FEB 1	998
WIPO	PCT	

A. The indications made below relate to the microorganism refer	red to in the description
on page13,	
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution	·
American Type Culture Collection	
Address of depositary institution (including postal code and count 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	(י <i>ר</i> י
Date of deposit	Accession Number
May 16, 1997	209054
C. ADDITIONAL INDICATIONS (leave blank if not applicable	(e) This information is continued on an additional sheet
EUROPE In respect of those designations in whi of the deposited microorganism will be the mention of the grant of the Europea application has been refused or withdraby the issue of such a sample to an exp the sample (Rule 28(4) EPC).  D. DESIGNATED STATES FOR WHICH INDICATIONS A	made available until the publication n patent or until the date on which wn or is deemed to be withdrawn, only ert nominated by the person requesting
E. SEPARATE FURNISHING OF INDICATIONS (leave blan	
The indications listed below will be submitted to the International Number of Deposit")	Bureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
This sheet was received with the international application	This sheet was received by the International Bureau on:
Authorized officer  Yaralagal Specialist  Faralagal Specialist  IAPD-PCT Operations  1APD-97-3746	Authorized officer

10

15

20

25

What is claimed is:

- 1. An isolated polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y; or a nucleotide sequence complementary to said isolated polynucleotide.
- 2. The polynucleotide of claim 1 wherein said polynucleotide comprises the nucleotide sequence contained in SEQ ID NO:X encoding the receptor polypeptide of SEQ ID NO:Y.
  - 3. The polynucleotide of claim 1 wherein said polynucleotide comprises a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length.
- 4. The polynucleotide of claim 3 which is polynucleotide of SEQ ID NO: X.
  - 5. The polynucleotide of claim 1 which is DNA or RNA.
- 6. A DNA or RNA molecule comprising an expression system, wherein said expression system is capable of producing a receptor polypeptide comprising an amino acid sequence, which has at least 80% identity with the polypeptide of SEQ ID NO:Y when said expression system is present in a compatible host cell.
  - 7. A host cell comprising the expression system of claim 6.
- 8. A process for producing a receptor polypeptide comprising culturing a host of claim 7 under conditions sufficient for the production of said polypeptide and
  30 recovering the polypeptide from the culture.
  - 9. A process for producing a cell which produces a receptor polypeptide thereof comprising transforming or transfecting a host cell with the expression system of claim 6 such that the host cell, under appropriate culture conditions, produces a receptor polypeptide.
  - 10. A receptor polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence of SEQ ID NO:Y over its entire length.

- 11. The polypeptide of claim 10 which comprises the amino acid sequence of SEQ ID NO:Y.
- 5 12. An antibody immunospecific for the receptor polypeptide of claim 10.
  - 13. A method for the treatment of a subject in need of enhanced activity or expression of receptor polypeptide of claim 10 comprising:
- (a) administering to the subject a therapeutically effective amount of an agonist to said receptor; and/or
  - (b) providing to the subject polynucleotide of claim 1 in a form so as to effect production of said receptor activity in vivo.
- 14. A method for the treatment of a subject having need to inhibit activity or expression of the receptor polypeptide of claim 10 comprising:
  - (a) administering to the subject a therapeutically effective amount of an antagonist to said receptor; and/or
  - (b) administering to the subject a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said receptor; and/or
  - (c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said receptor for its ligand.
  - 15. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of the receptor polypeptide of claim 10 in a subject comprising:
  - (a) determining the presence or absence of a mutation in the nucleotide sequence encoding said receptor polypeptide in the genome of said subject; and/or
- (b) analyzing for the presence or amount of the receptor polypeptide
   30 expression in a sample derived from said subject.
  - 16. A method for identifying agonists to the receptor polypeptide of claim 10 comprising:
    - (a) contacting cells produced by claim 9 with a candidate compound;
- 35 and

25

(b) determining whether the candidate compound effects a signal generated by activation of the receptor polypeptide.

10

- 17. An agonist identified by the method of claim 16.
- 18. The method for identifying antagonists to the receptor polypeptide of claim 10 comprising:
  - (a) contacting said cell produced by claim 9 with an agonist; and
- (b) determining whether the signal generated by said agonist is diminished in the presence of a candidate compound.
  - 19. An antagonist identified by the method of claim 18.
- 20. An isolated receptor polynucleotide comprising a nucleotide sequence selected from the group consisting of:
  - (a) a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC; and
  - (b) a nucleotide sequence complementary to the nucleotide sequence of (a).
- 21. A recombinant host cell produced by a method of Claim 9 or a membrane thereof expressing a receptor polypeptide.

	10 20	30
	MLLLLLLPLLWGRERVEGOKSNRKDYSL WLPLLLPLLWG GSLOEKPVYEL	GOKSNRKDYSLTMOSSVTVOEGM HMACR70.AA GSLOEKPVYELOVOKSVTVOEGLOB-1.aa
<b>4</b> %	CVHVRCSFSYPVDSQTDSDPVHGYWFRACVLVPCSFSYPWRSWYSSPPLYVYWFRD	FOUNDISWKAPVAT HMCR70.AAPLYVYWFR DGEIPYYAEVVAT OB-1.aa
88 92	NNPAWAVQEETRDRFHLLGDFQTKNCTLSIRDARMSDAGR NNPDRRVKPETQGRFRLLGDVQKKNCSLSIGDARMEDTGS	110 120 SIRDARMSDAGR HMCR70.AA SIGDARMEDTGS OB-1.aa
121	YFFR MEKG-NIKWNYKYDOLSVYFFR VERGRDVKYSYQQNKLNLEVTALI	150 160 HMCR70.AA EKPDIHFLEPLE OB-1.aa
142 156	170 180 190 	190 200 

F16.1A

	P HMACR70.AA A OB-1.aa	7 280 L D HMACR70.AA P OB-1.aa	110 Q - V H L G D E G E F HMACR70.AA R R V R S A E E G G F OB-1.aa	HMACR70.AA OB-1.aa	HMACR70.AA OB-1.aa
240	T Y P	280 A V D D A P	320 B B C B		400  P W A
	N V T O L N V S	RLVC	L G D S A E	- A - E - C	S S S A G
230	ERTV	270 LSVLEGQSLRLVCAV LPVLEGQALRLLCDA	310 LELQ-VHLGD LELRRVRSAE	350 	390  F K V N S
	TTAO	N G G G G G G G G G G G G G G G G G G G		1 G P S I S	1 S I S
	R Q G A	S E S S E S E S E S E S E S E S E S E S	S S S S S S S S S S S S S S S S S S S		880 
220	NVTYP GTNLTCQMKRQGAQVTTERTVQLNVSYA	260 GTASTALGNSSS GIALEILONTSY	300 I Y P S Q P S N P L V I N A T P I S N T G I	LNLS VYSLPQLLGPSCSWEAEGLHCR	380  KPLE
	T	TAPLE	SLTIYSPAIN	VSLNLS	- R - I - E - E
210	•	250 QGE R-NG	3 Q R D	1	370  S L C W
	- T - A - A	T V T	290 RLSWTWR HLSWFQG	S L G H P L G	3  RARPAP
	SELTLTPRPEDH	PQNLTV PQQq	SNPPARLSWTSNPPAHLSWF	330 TCRAQNSLGSQH TCRAQHPLGFLQ	SFRA
	142 - 196 S	147 P 236 P	187 S 273 S	226 T 313 T	244 - 353 C :

F.C. 18

HMACR70.AA OB-1.aa	HMPCR70.AA OB-1.aa	520 - M HWCR70.AA P L OB-1.aa	HMACR70.AA OB-1.aa
	480 A R P HMACR70.AA A G R OB-1.aa	520 M HAPP	
LOGR	R K K K K A K A K A K A K A K A K A K A	P G D A I	B B I K T S
430 1	470 V V R S C I V K A R	510  D Q A S P	550  ITEYS
	4. IF FLI	1 S C C	I   I   I   I   I   I   I   I   I   I
420 430 430 440 KMKPV	460 LSFCV ICLCL	500 510 520 M HMACR70 RKKPWPDSPGDQASPPGDAPPL OB-1.aa	540 
K P V Y X V S C K	GATALVF GVMALLC	T I I S G S R	540 S G . S F M K S R E P K D O E A P S T T E Y S E I K T S K
			는 다
410 LQQEYTG SSLILHGGLSS	450 LGAVG PAALG	490 D I G D E D P I	S30
N - N - N - N	450 - S G V L L G A V G G T G T G V V P A A L G G A	490 A A D V G D I G P E K M D D E D P I M G	530 KDANTIQ GLS
244 - 393 1	256 - 433 (	294 7	303 1

FIG. 10

1 H T P Y S S L K K L HTEDK48.aa S S L K L L MRC-OX44.aa A G L A V M PETA-3.aa	N G F V A V S G I I HTETK48.aa V Q N T Y G I L MRC-0X44.aa L K S D Y I S L PETA-3.aa	110 1 L H V G N L C L V M G C I T HTETK48.aa I M V V MRC-OX44.aa V M V T PETR-3.aa	LIMEVTAATVV HEDK48.aa LLAEVTLAILL MRC-OX44.aa FLLEIIAGILA PETA-3.aa
20 	60 SL GCCILGFGIBL GCCILGFGIHLL AVGIWTLA	100 L G L S S A Y L V U I V G S I L V V A G T V	140 LLFCILSMVIV LMSFFVLLLLI LRLYFILLLI
10 MAE MGM	50 I X Y V L F F F N F L F W V C G C C I	ENGEGIGGKCGGASLTNV FRNLPF LTLGNV LASGTY LATAYI	VLLGCAGWYGATKESRGT AFLGCMGSIKENKCL GVLGCCATFKERRNL
. ——	. 15 . 36 . 36	8 <del>2</del> 23	 69 75

-16.2A

		<i>4 1</i>	
200 W N HTEDK48.aa W D MRC-OX44.aa V D PETA-3.aa	240 G H HTEDK48.aa MRC-OX44.aa PETA-3.aa	280 T K HTELK48.aa F H MRC-OX44.aa I Q PETA-3.aa	320 L L HTEDK48.aa A L MRC-OX44.aa L E PETA-3.aa
190 NY-RGYNEPDDYSTQW SI-QHYHSDNSTRMAW TMTKRYHQPGHEAVTSAV	WIRSQEAG	PNVIHOKO PSGADVO SNIYKVE	310 RYV A Q A G L E SFA LTLNCQIDKTSQ IFTCC LYRSLK
170 180 L L F F P I V G D V A L E H T F V T L R K F V Y E K K I N T L V A E G L N D Y A Y Y Q Q L N T E L K E N L K D	LVMEKLKCO FIQSQLQCO QLQQEFHCO	T Y P R S C C K S P D S C C K T	290 300 300 1
109 104 112	146 137 148	174 158 181	213 180 218


FIG.2B

MRC-0X44.aa HTEDK48.aa PETA-3.aa 245 218 252

FIG. 2C

HPW£E25.aa NAG-2.aa HPWÆE25.aa NAG-2.aa TALLA-1.aa HPWAE25.aa NAG-2.aa TALL≏-1.aa **<** > A IT (2 G G V - 5 Ŧ Þ  $\times \times \sigma$ <u>о</u> а <u>п п о</u> A SSI  $\forall \forall \vdash$ ပ ပ 5 5 4 338 38 12 12 1 10 00 00

130 140 150 150 160 160 150 160 160 150 160 160 170 W N T T M K G L K C C HPWAE25.aa 113 R Y A Q Q D L K K G L H L Y G T Q G N V G L T N A W S I I Q T D F R C C NAG-2.aa 113 D T F L R T Y T D A M Q T Y N G N D E R S R A V D H V Q R S L S C TALLA-1.aa	170 200 51 GFTNYID - FEDSPYFKENSAFPPFCCNDDVTNT : NETCTK HPWAE25.aa 49 GVSNYIDWFEV YNATRV - PDSCCLEF SESCGL NAG-2.aa 47 GVQNYIN - WSTSPYFLEHGIL - PPSCCMNE TD - CNP TALLA-1.aa	210 220 230 240 90 QKAHDQKVE G G F N Q L L Y D I R T N A V T V 3 G HPWAE25.aa 80 H A P G T W W K A P G Y E T V K V W L Q E N L L A V 3 I F G L C T NAG-2.aa 79 Q D L H N L T V A A T K V N Q K G G Y D L V T S F M E T N M G I I : G TALLA-1.aa	250 270 280 18 VAAGIG-GLELA-AMNOVHVSVLQSTISPLLPLPLLL HPWAE25.aa 13 ALVQIL-GLTFAMTMYONAG-2.aa 14 VAFGIAFSQLIGMLLACCLSTALLA-1.aa	290 300 900 900 900 900 900 900 900 900 9	FIG.3B
110	151 149 147	190 180 179	218 213 214	253 229 234	




HG.4A

		<i>7121</i>	
HTPEF86.aa B1.aa	HIPEF86.aa Bl.aa	HTPEF86.aa Bl.aa	HTPEF86.aa B1.aa
240 EFGI	280 N PW I T P E E E W V G L T	320  QDQESSP	
220 230 230 240 1 S G V L L V F C L L E F G I 1 L S V M L I F A F F Q E L V I A G I V E N	270 1 Y P N I Y A A 1 E I K	310 320  ETETNFPEPPQDQESSP	
220   NA I S G V L L V   G I L S V M L I	Z60 Z70 Z80 LVCCQSSNVSVIYPNIYAANPWITPE LLSAEEKKEOTI - EI KEEVVGLT	300  PIQEEBEBGIG	
210 MA SPSTQYCYSIQSLFLG	250 ACASSHFGCQLV EWKRTCSRPKSNIVLL	290 S P P S Y S S E I Q S Q P K N E E D I È I I	ANK. IENDSP
182 175 K N	199 215 EWI	235 PVT 251 ETS	248
<del>~</del> ~	H 73	0.0	Ø Ø

FIG. 4B

ις 1 Γυ 1	S R R L Q	T K P V I T K P V I T	20 20 FCFKSVLL FCLKTLLI	I X I X	TEIFWITGVILLAVGI SFVFWITGVILLAVGI	LAVGI LAVGV	HSBBF02.aa TALLA-1.aa
S E	LEENY	FSLUNE	E K A T N V E N S T N A		EE	_	HSBBF02.aa TALLA-1.aa
R A S A S S S S S S S S S S S S S S S S	Z Z Z	A W M L K L Y A M F P W M L K L Y A M F	TOOT TOO TOO TOO TOO TOO TOO TOO TOO TO	FLTLVFLVELVAAIVGFV FLSLVFLAELVAGISGFV	AIVGFVFR GISGFVFR	HEH	K D TALLA-1.aa
N X X X Y	XU	ALKOYNST AMOTYNGN	<u>П</u> <u>О</u>	Y R S H A V D K I Q N T - R S R A V D H V Q R S	ONT LHCCGV	V T D Y R V O N Y T	TDYR HSBBF02.aa ONYT TALLA-1.aa
N A A	170 Y Y S E	K G F P H G I P	TE CCKL	E - DCTPQ ETDCNPQ	180 190 200 C C K L E - D C T P Q R D A D K V N C C M N E T D C N P Q D L H N L T V A A T K V N		HSBBF02.aa TALLA-1.aa


F16.5A



SUBSTITUTE SHEET (RULE 26)


		•		
HLTAH80.aa TALLA-1.aa	HLTAH80.aa TALLA-1.aa	80.aa -1.aa	30.aa -1.aa	30.aa -1.aa
нглан тагта	HLTAH TALLA	HLTAH80.aa TALLA-1.aa	HITAH80.aa TALLA-1.aa	180 200 RETASNONGSLA-HPSDLYA HITAHBO.aa MNET-DCNPODLHNLTVAATKVNQ TALIA-1.aa
	- ജ - യ .ഗ	-84+x		20- 40- 40-
E E	X X	R R I I Z	160 160 WENT WSTS	L Y
F F I	E E	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	ΩZ	N E
Z K	GCCATIREGCFATCRG	VENEVDIKDTEL	S K	H F
0 A	<b>5</b> 5	D H W	T H N X	A T
30 40 1 T Y D T T T T T T T T T T T T T T T T T	70 GALLFIIGLI GTTIVVEGLF	110 R A K R H E	HCCGIHNYSDWENTSCCGVQNYINWSTS	190 S N
11 12	H F F	> > Y F ¹	VOROLHCCG VORSLSCCG	O
& B C B	7 H	₩ ₩	0 0	10
FWGAAG FWITGV	T T T	VVVL GY AGISGF	2 R	Z Z
Z Z Li	<u>ი</u> ი	> &   		N Q
- 8-H >	- 8-4 6 4 6 7 H	-0->>	140 A I D Y A V D H	180 T B T
V F L N L I I I Y S F V	V I	V T E V L A E L	SRATSRAN	M M H N
→ H H	A V X	<u> </u>	A S S	υ υ
		<u>1</u> 1	DDE	17 CA 10 CA
K T V K T L	T S T F	1 J	N P	<b>₽</b> ₽
S - 10 C S - 10	> z	ж н н ж	130 N G T N G T	170 H G
E E	E D	TFVI	T Y N G	К N Q
C G	E H	A X	X O	<u>E</u> 1
H H	D H I S	M L K K	ж <u>ү</u> х	는 도 보기
ច ២	T X	Ж Ф О 🔀	OH	D W
<del>디</del> 디	8 4	88	120	160 158

16.6A



		14/2/		
HTPBA27.aa NAG-2.aa	HTPBA27.aa NAG-2.aa	HTPBA27.aa NAG-2.aa	HTPBA27.aa NAG-2.aa	HTPBA27.aa NPG-2.aa
FNLFFWLGGCGVLGVGIWLAATQGS FNLLFWLGGCGVLGVGIWLAATQGS	60 70 80 NLLILTGAFVMAIGFVGCLGAIKENK	110 120 FAYTDKIDRYAQQDLK FAYTDKIDRYAQQDLK	TNAWSIIOTDFRCCGVSNYTDWFEVY	ESCGLHAPGTWWKAPCYETVKVWLQE
AA	60 ANLLITGAFVMA ANLLITGAFVMA	100 LLEATIATLF LLEATIAILF	140 I TNAWSIIQTDFR I TNAWSIIQTDFR	
10 MARACLQAVKYLMF MARACLQAVKYLMF	FATLSSSFPSLSAA FATLSSSFPSLSAA	CLLLTFFLLLLLVF CLLLTFFLLLLLLVF	KGLHLYGTQGNVGL KGLHLYGTQGNVGL	170 NATRVPDSCCLEFS NATRVPDSCCLEFS
	च च	# # # # # # # # # # # # # # # # # # #	121	161 161

FIG./A



F16.8A

CD9 antigen.aa CD9 antigen.aa CD9 antigen.aa HAIDO59. aa IGIHADD59.aa IGA CD9 antiger HAIDQ59. aa თ ი ი 160 200 回 A 0 0 7 7 LOL FRI TIISVEVEDN X T A A I 150 230 190 田区 口 日 D B B ZK 民民 ÖЫ C X X S 1. 24 CAC DRGKGN TKDEPOR 140 D T V 180 220 **ပ** M V IS 니도 고 E च च S CPK----CPKKDVL I W M I N K GLTIFGN VVMIFGN A Y N T Y N 130 170 210 田口 EH 田式 O 0 Ŀı 9 1 0 g OШ **B** > 923 158 159 192 199

FIG.8B

æt	 mt		
.40. <i>a</i> .	140.az 13.aa	:40.až :3.aa	.40. <i>až</i> 3.aa
HHFEK40.aa PEIR-3.aa	STFAASAYILIFAGVLVMVTGFLGFG HHFEK40.aa GTYLATAYILVVAGTVVMVTGCC PEPA-3.aa	HHFEK40.aa PETA-3.aa	HHFEK40.aa PEIA-3.aa
	- & <u>- o o</u>		- 9- 5 E
G H	<u>Б</u> О	H H	S S N
A V A V	T N L L	0 0 0	ບ. ບ <b>ບ</b> ບ
Z Z	O,D	X X	O O
A	I V	<u>&gt;</u>	E E
ט ט	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	A A	0,0
- 유 - 영 - F	1 C L - S -	11- 11- 11- 11-	- 1 - 0 - 0 - 0
F F	A G.	A G	B: D
F C	I F V V	LVII	> >
Z Z K H	11 11	<u> </u>	A H
> E4	XX	13 12	- C
- 8- 11	- 8- n F	-11-10	140 E A L
Y 1	A J		回 Q
Y L K Y L, E, F V F N F F F W V G G A A V L A V G I W C L K Y L, E, F T Y N C C F W L A G L A V M A V G I W		100 110 12C LLLVIFLVELVAGVLA HVYYQRLSD LLLIIFLLEIIAGILA YAYYQQLNT	140
C K	S S	O H	H K
1 D	L A L A	K K L H	l z z
5 0	- 52 - 02 (0  - 12	\\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\	130 LA E M T K
ΔE	니니	- 영- 타디	
E X O X	K D C	z z	N N U
X Z F H	$x \times x$ $x \times x$	स्य स्य स्य	H Z
X H E F	의 고 > 전	XX HH	ス ス 〇 円
E E	H H	A A	ы ы ы
	4 4	ස්ස්	121

FIG.9A





FIG.9B

			20 -	30.	
ਜਜ	MCTGKCARC MCTGKCARC	CVGLSLITLCEVC	V C I	VANALLLVPNGETSWTNT VANALLLVPNGETSWTNT	HGBGV89.aa L6H.aa
4 4	NHLSLQVWLMGG NHLSLQVWLMGG	<b>H</b> H	GGGLMVLCPGIAAVE	GGGLMVLCPGIAAVRAGGKGCCGAGC	1 80 - C HGBGV89.aa C L6H.aa
. සස	GGNRCRMLRS	VFSS VFSS	нн	CC	120   HGBGV89.aa   L6H.aa
121	NGEWGYHFE NGEWGYHFE	130 DTAGAYLLNR DTAGAYLLNR	140 Y L L N R T L W D R C E A P E Y L L N R T L W D R C E A P E	CEAPPRVVPWNVTLFS	0 HGBGV89.aa L6H.aa
161 161	LLVAASCLE LLVAASCLE	170 180 IVLCGIQLVNAT IVLCGIQLVNAT	I G V F C G D I G V F C G D	190 CRKKODTPH CRKKODTPH	HGBGV89.aa L6H.aa

FIG. 10

	10				<b>-</b> 4-
の 24 ひ ひ 至 22	RKCGGGLS GKCARCIG	CLLIPLALWS HSLVGLALLC	WSIIVNILLYFPNG LCIAANILLYFPNG	IPLALWSIIVNILLYFPNGOTSYAS VGLALLCIAANILLYFPNGETKYAS	S S HUVBB80.aa S E L6.aa
NN	SOTINY VWYFE	GICFSGIMML GIVGGGLLML	60 NL IVTTVLL NL LBAFVFI	T VTTVLLVLE NNNYKÇCO LBAFVFIGEEODDCGGCCG	80 1 Q HUVBB80.aa G L6.aa
N N E E N N	SKKYVTL CGKRCAML	LSIIFSSLG SSVLAALIG	100 GIAFSGYCL GIAGSGYCV	100 GIAFSGYCLVISALGLVQGPY GIAGSGYCVIVAALGLAEGPL	120 Y HUVBB80.aa L L6.aa
0 0 - 1 D R	T.L.D.G.W.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.A.E.Y.E.Y	FEGTAGRE FASTEGOY	140 L T D S S I W L L D T S T W	150 160 160 160 160 1 S E C T E P K H I V E W N V	160   I HUVBB80.aa   V L6.aa

FIG. 11A



HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HJACE54.aa rGalectin-5.aa hGalectin-8.aa
40 G L S P G Q V N G L Y P S K S D O L D P G T L	80 	120  EIVIMVL	160  S I G
30 VPCSHALP VPFFTSIPN IPFVGTIP	70   MKPRADVAF	110 	150  DTLGIYGKVNIH
20 	V S L R D Q A A I N L R C - G G V D L Q N - G S S N	100 	140 
10 	SO VRGLVLQEPKHFT ISGVVLSDAKRFQ IRGHVPSDADRFO	90 	130  KFQVAVNGKHTLLY
2 2 Z	23 33 34 14 14 14 14 14 14 14 14 14 14 14 14 14	25 R R R R	46 53 112 K A

FIG. 12A

	170	180	190	200 HJACE54.aa
1 03 1	SFSDLQSTQAS	SLELTEISRE 220	NVPKSGTPQLR 1	LPFARR Malectin-5.aa LPFAAR Malectin-8.aa
			H D F N V D L L A G K S K D	APVT IAFH IALH
	RASFADRTLAWIS NPRFDENAVVRNT NPRLNIKAFVRNS	260 RWG-QK QINNSWGPEE FLQESWG-EE	Z70 KLISAPFLFYP RSLPGSMPFSR ERNITSFPFSP	280 QRFEVHTACE54.aa GQRFSVACAlectin-5.aa GMYFEMHGALectin-8.aa
	L L L F Q E G G L K L A L N W I L C E G H C F K V A V D I I Y C D V R E F K V A V N	300 G Q G L G A T S M G Q H I C E Y S H G V H S L E Y K H	310 N Q Q A L E Q L R E L R L M N L P D I N T L R F K E L S S I D T L	RISGSV HJACE54.aa EVAGDI rGalectin-5.aa EINGDI hGalectin-8.aa
	330 Y C V H S . T H V E T L E V R S W	FIG.12B	28	HJACE54.aa rGalectin-5.aa hGalectin-8.aa




FIG. 15

cited i Repor Your f	n the Europear t of EP <i>00 Q2</i> Ref.: PF 3.54 PC	Search 35/27 35365 73 EP	<b>7</b>	
HMMSS46.aa B-cell Recpt Associated Pr.aa	HMN5S46.aa B-cell Recpt Associated Pr.aa	i.aa Recpt Associated Pr.aa	HMMSS46.aa B-cell Recpt Associated Pr.aa	Associated Pr.aa
		HMMGS46 B-cell		HMMSS46.aa B-cell Recpt Associated
MAQNLKDLAGRLPAGPRGMGTALKLLLGAG	AVAYGVRESVFTVEGGHRAIFFNRIGGVQQ	1 DTILAEGLHFRIPWFQYPIIYDIRARPRKI DTILAEGLHFRIPWFQYPIIYDIRARPRKI	100 110 120 1 SSPTGSKDLQMVNISLRVLSRPNAQELPSM 1 SSPTGSKDLQMVNISLRVLSRPNAQELPSM	YORLGLDYEERVLPSIVNEVLKSVVAKFNA
ਂ ਜ ਜ	ਸ਼ ਸ਼	ਰ ਰ	<b>ਲੇ</b> ਲੋ	22 22

FIG. 14A

HMMSS46.aa B-cell Recpt Associated Pr.aa HMMSS46.aa B-cell Recpt Associated Pr.aa	B-cell Recpt Associated Pr.aa  HWMGS46.aa  B-cell Recpt Associated Pr.aa  HWMGS46.aa  B-cell Recpt Associated Pr.aa
160 170 SQLITQRAQVSLLIRRELTERA SQLITQRAQVSLLIRRELTERA 190 200 VAITELSFSREYTAAVEAKQV VAITELSFSREYTAAVEAKQV  7220 230 FLVEKAKQEQRQKIVQAEGEA	OFEVERAROEOROKIVOAEGEAEAAKMLGE  250 260 270 ALSKNPGYIKLRRIRAAONISKTIATSONR ALSKNPGYIKLRRIRAAONISKTIATSONR IYLTADNLVLNLODESFTRGSDSLIKGKK

FIG. 14B

30 FMH HNFGWD6.aa	60 G M A A HNFGAD6.aa EGFR related-protein.aa	90 L PIP HWFGWD6.aa EGFR related-protein.aa	120 C L F HNFGWD6.aa EGFR related-protein.aa	150 V P G HNFGAD6.aa EGFR related-protein.aa	HNFGWO6.aa EGFR related-protein.aa
ZO TKGSCEITTREYCE TKGRCEITSREYCD	SO RRGRPGVVEERTLG H	80 ASDSGCFWGAEHHM	110 FLNPEVPDOFYRSG FLNPEVPDOFYRL-	140 PTPVMDTQADPWGK W	PEVSG AGILHC FIG.15
MDCEIKGRPCCIG	GYFHEEATLCSOV GYFHEEATLCSOV	70 CWGRGSRTPSHVG	100 RCTVLDKVCWAAA CMDDVCGLLP	130 SYMLGKRSSMPPN 	160 PGYGRSNEPKTTA
<del></del>	표표	ය 3	ደ ዓ	121	151 69

International Application No PCT/US 98/00959

		<del></del>	
A. CLASSI IPC 6	C12N15/12 C12N15/85 C12N5/1 C12Q1/68 A61K38/17 A61K48/		C07K16/28
According t	o International Patent Classification (IPC) or to both national classific	ation and IPC	
B. FIELDS	SEARCHED	•	
Minimum de IPC 6	ccumentation searched (classification system followed by classification C12N C07K C12Q A61K	on symbols)	
	tion searched other than minimum documentation to the extent that s		
Electronic d	ata base consulted during the international search (name of data ba	se and, where practical, search te	rms used)
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the rel	event passages	Relevant to claim No.
X	DATABASE GENBANK Accession No. T48852, 8 February "EST70319 from Stratagene place XP002066898 see the whole document		1-12,20
X	DATABASE GENBANK Accession No. G14442, 4 January "STS SHGC-10055" XP000206689 see the whole document	1996	1-3,20
		-/	
[V] =	ner documents are listed in the continuation of box C.	Patent family members a	era listad in anna-
<u> </u>		Patent lamily members	ue nated in criter.
*Special categories of cited documents:  "A" document defining the general state of the art which is not considered to be of particular relevance  "A" document defining the general state of the art which is not considered to be of particular relevance  "I later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention			
"E" earlier of filing d	ocument but published on or after the international ate	"X" document of particular releval	
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  "L" document which may throw doubts on priority claim(s) or involve an inventive step when the document is taken alone 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the			
other r "P" docume	"O" document referring to an oral disclosure, use, exhibition or other means document is combined with one or more other such document of the means such combination being obvious to a person skilled in the art.    International fitting date but   International fit		
Date of the	actual completion of the international search	Date of mailing of the internal	<u> </u>
4	June 1998	_	9. 1998
Name and n	nailing address of the ISA  European Patent Office, P.B. 5818 Patentiaan 2	Authorized officer	
NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo ni, Fax: (+31-70) 340-3016  Galli, I			

Internatic—al Application No
PCT/US 98/00959

<u> </u>		PC1/03 98/00959
C.(Continua	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	Relevant to claim No.
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<b>A</b>	ZANNETTINO A.C.W. ET AL.: "A powerful technique for isolating genes encoding surface antigens using retroviral expression cloning" J. IMMUNOL., vol. 156, no. 2, 15 January 1996, pages 611-620, XP002066893 see abstract	1-21
<b>A</b>	ADAMS M D ET AL: "COMPEMENTARY DNA SEQUENCING: EXPRESSED SEQUENCE TAGS AND HUMAN GENOME PROJECT" SCIENCE, vol. 252, no. 5013, 21 June 1991, pages 1651-1656, XP000645049 cited in the application see abstract	1-21
A	SIMMONS D. & SEED B.: "Isolation of a cDNA encoding CD33, a differentiation antigen of myeloid progenitor cells" J. IMMUNOLOGY, vol. 141, no. 8, 15 October 1988, pages 2797-2800, XP002066894 see abstract see figure 4	1-21
<b>A</b>	CROCKER P.R. ET AL.: "Sialoadhesin, a macrophage sialic acid binding receptor for haemopoietic cells with 17 immunoglobulin-like domains" EMBO J., vol. 13, no. 19, 3 October 1994, pages 4490-4503, XP002066895 see abstract see figure 2	1-21
A	KELM S. ET AL.: "The Sialoadhesins: a family of sialic acid-dependent cellular recognition molecules within the immunoglobulin superfamily" GLYCOCONJ. J., vol. 13, no. 6, December 1996, pages 913-926, XP002066896 see the whole document	1-21
A	DATABASE GENBANK Accession No. U71382, 12 November 1996 PATEL N. ET AL.: "OB binding protein - 1" XP002066901 cited in the application see the whole document	1-21

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

International Application No PCT/US 98/00959

•		PC1/05 98/00959
C.(Continue Category *	etion) DOCUMENTS CONSIDERED TO BE RELEVANT  Citation of document, with indication; where appropriate, of the relevant passages	Relevant to claim No.
Calegory	Charles of accountant, with transmission appropriate, of the teneval passages	
P,X	DATABASE GENBANK Accession No. AA344713, 21 April 1997 "EST50650, H. sapiens cDNA similar to CD33 antigen" XP002066900 see the whole document	1-12,20
P,A	TAKEI Y. ET AL.: "Molecular cloning of a novel gene similar to myeloid antigen CD33 and its specific expression in placenta" CYTOGENET. CELL GENET., vol. 78, 1997, pages 295-300, XP002066897 see abstract see figure 1	1-21
	-	
•		
	•	
	,	

Inter onal application No.

PCT/US 98/00959

Box I	Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)
This Inter	rnational Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely: Remark: Although claims 13 and 14 are directed to a method of treatment of the human/animal body, and claim 15 (as far as in vivo methods are envisaged) is directed to a diagnostic method, the search has been carried out and based on the alleged effects of the compound/composition.  Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3.	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II	Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)
	ernational Searching Authority found multiple inventions in this international application, as follows:
1.	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee
3.	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. X	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  see further information sheet, subject 1.
Remari	The additional search fees were accompanied by the applicant's protest.  No protest accompanied the payment of additional search fees.

何のできる。最終に対けなった。 きこれをはが過ぎて大阪のでは、古典をのははな

#### FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: (1-21) - partial

An isolated polynucleotide encoding the HMACR70 cell surface receptor (Seq. IDs 1,18), polynucleotides complementary thereto, fragments thereof, homologs thereof.

An expression system comprising said polynucleotides, a process to produce a cell harbouring said expression system. Said cell or membrane thereof expressing said receptor.

A process for the production of the HMACR70 polypeptide comprising culturing said cell.

Said HMACR70 polypeptide or homologs. An antibody directed against said polypeptide.

Therapeutic and diagnostic applications.

A method to identify agonists or antagonists to said receptors. Said agonists and antagonists.

2. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 2,3,19

3. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 4,20

4. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 5,21

5. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 6,22

6. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 7,23

7. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 8,24

### FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

- 8. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 9,10,25
- 9. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 11,26
- 10. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 12,27
- 11. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 13,28
- 12. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 14,29
- 13. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 15,30
- 14. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 16,31
- 15. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 17,21

# This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

BLACK BORDERS

IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

FADED TEXT OR DRAWING

BLURRED OR ILLEGIBLE TEXT OR DRAWING

SKEWED/SLANTED IMAGES

COLOR OR BLACK AND WHITE PHOTOGRAPHS

GRAY SCALE DOCUMENTS

LINES OR MARKS ON ORIGINAL DOCUMENT

REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

## IMAGES ARE BEST AVAILABLE COPY.

OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.